

## THE ROLE OF *DREB2* GENE IN DROUGHT TOLERANCE OF COMMON WHEAT (*TRITICUM AESTIVUM* L.) ASSOCIATED WITH *AZOSPIRILLUM BRASILENSE*

DEHGHANI, I.<sup>1,5</sup> – MOSTAJERAN, A.<sup>2\*</sup> – ESMAEILI, A.<sup>3</sup> – GHANNADIAN, M.<sup>4,5</sup>

<sup>1</sup>University of Isfahan, Plant Science Division, Department of Biology, Faculty of Science  
Isfahan, Iran  
(phone: +98-91-39041386; fax: +98-31-37932456)

<sup>2</sup>University of Isfahan, Plant Science Division, Department of Biology, Faculty of Science  
Isfahan, Iran  
(phone: +98-31-37932471; fax: +98-31-37932456)

<sup>3</sup>University of Isfahan, Cell, Molecular Biology and Biochemistry Division  
Department of Biology, Faculty of Sciences, Isfahan, Iran  
(phone: +98-31-37932452; fax: +98-31-37932456)

<sup>4</sup>Isfahan University of Medical Sciences, Isfahan Pharmaceutical Sciences Research Center  
School of Pharmacy, Isfahan, Iran  
(phone: +98-31-37921335; fax: +98-31-36680048)

<sup>5</sup>Isfahan University of Medical Sciences, Department of Pharmacognosy, School of Pharmacy  
Isfahan, Iran  
(phone: +98-31-37921335; fax: +98-31-36680048)

\*Corresponding author

e-mail: mostajerana@yahoo.com; phone: +98-31-37932471; fax: +98-31-37932456

(Received 31<sup>st</sup> Jul 2018; accepted 27<sup>th</sup> Feb 2019)

**Abstract.** Dehydration causes loss of wheat (*Triticum aestivum* L.) yield. Inoculation of wheat with *Azospirillum brasilense* improves its tolerance to drought. Although up-regulation of *DREB2* gene has increased drought tolerance of wheat, less information exists about *DREB2* expression under inoculation with *A. brasilense*. In this study, the physiological responses of different pairs of six wheat cultivars with *A. brasilense* Sp7 and Sp245 strains were evaluated to drought and the tolerant (Roshan-Sp245), sensitive (Shahpasand-Sp7), moderately tolerant (Roshan-Sp7) and moderately sensitive pairs (Shahpasand-Sp245) were selected. Afterward, in the second experiment, *DREB2* expression of the selected pairs was evaluated under inoculation and/or dehydration ( $\psi_w = -90$  MPa) at 0, 120 and 360 minutes. At 120 min, *DREB2* expression was more in the tolerant and moderately tolerant pairs than others and also higher in inoculated and/or dehydration conditions. Root's *DREB2* was up-regulated almost 700-fold in the tolerant and moderately tolerant pairs under dehydration. In contrast, in the sensitive cv., *DREB2* expression did not change due to dehydration or inoculation. This could be the effect of compatibility or homology of *A. brasilense* strains and wheat cultivars which affected *DREB2* expression. Therefore, *DREB2* expression had a significant effect on increased drought tolerance in tolerant and moderately tolerant pairs.

**Keywords:** wheat, *Azospirillum brasilense*, Sp7, Sp245, drought tolerance, *DREB2* gene, dehydration

**Abbreviations:** *DREB2* gene, DRE-binding factor gene; FC, Field capacity; PAL, Phenylalanine Ammonia lyase enzyme (EC 4.3.1.24); TAL, Tyrosine Ammonia lyase enzyme (EC 4.3.1.23); RDW, Root Dry Weight; SDW, Shoot Dry Weight; MSL, Maximum Shoot Length; ARL, Average Root Length; NMR, Number of Main Roots; NRB, Number of Root Branches; RTP, Root Total Protein; STP, Shoot Total Protein; RPAL, Root PAL activity; SPAL, Shoot PAL activity; RTAL, Root TAL activity; STAL, Shoot TAL activity

## Introduction

Drought is one of the major or more significant stresses which adversely affect the plant water status and crop yield (Kramer and Boyer, 1995; Bray, 2000). Dehydration or loss of water causes reduction of cell turgor pressure, an increase in negative water potential and consequently less plant production (Liu et al., 2005). *Azospirillum brasilense* is one of the PGPRs (Plant Growth Promoting Rhizobacteria) that can associate with wheat and increase crop quality and quantity through suitable physiological and biochemical modification in plants (Dobbelaere et al., 2003; El-Samad and Hetta, 2005; Pereyra et al., 2006; Hungria et al., 2010; Veresoglou and Meneses, 2010). The effect of PGPRs on dehydration responsive genes in plants has been evaluated but not well studied specifically about the *DREB2* gene expression during inoculation of wheat with *A. brasilense*. Therefore, the objective of this study is to evaluate the effect of *A. brasilense* on *DREB2* gene expression in wheat cultivars under dehydration condition.

## Review of Literature

The ability of *Azospirillum* to establish an association system with plants such as wheat through which it promotes better growth and development (Dobereiner and Day, 1976; Bashan and Holguin, 1997) is due to its capacity for N<sub>2</sub> fixation capacity (Bermner et al., 1995) and production of some bacterial compounds having plant phytohormone properties such as nitric oxide, IAA (indole-3-acetic acid) polyamines and amino acids (Hartmann et al., 1994; El-Khawas and Adachi, 1999; Pakdaman et al., 2014). These bacterial compounds produced in wheat rhizosphere caused structural changes in wheat roots, stimulated root growth and increased water and minerals absorption by the plant (Perrig et al., 2007; Amooaghaie et al., 2004; Casimiro et al., 2003; Correa-Aragunde et al., 2004). These advantages for plants are also reported at dehydration condition for different pairs of plant and bacteria especially at the early stage of growth (Bashan and Holguin, 1997; Hartmann and Baldani, 2006; Yang et al., 2009). For instance, promotion of dehydration tolerance is reported in different associated pairs obtained from some *A. brasilense* strains and wheat (*Triticum aestivum* L.) cultivars (Díaz-Zorita and Fernández-Canigia, 2009).

PGPRs can affect the expression of stress-responsive genes and thus modulate plant responses to stress (Jakab et al., 2005; Gachomo et al., 2014; Barnawal et al., 2017). For example, up-regulation of *RD29A* gene in *Arabidopsis* plants inoculated with a PGPR strain (*Bradyrhizobium japonicum* IRAT FA3) was observed when compared to non-inoculated plants (Jakab et al., 2005; Gachomo et al., 2014). *DRE/CRT* (Dehydration responsive element/C-repeat), a cis-acting element in the promoter region of *RD29A* (Yamaguchi-Shinozaki and Shinozaki, 2005; Jia et al., 2012; Gachomo et al., 2014) interacts with a series of transcription factors known as DREBs proteins (DREB/CBF; DRE-binding factor/C-repeat binding factor) inducing a set of downstream dehydration responsive genes that improve dehydration tolerance of plants (Baker et al., 1994; Sakuma et al., 2002; Yamaguchi-Shinozaki and Shinozaki, 2005). *DREBs* genes family which encode DREBs proteins has a main role in the gene expression and regulation of plant's response to dehydration stress (Sakuma et al., 2002; Yamaguchi-Shinozaki and Shinozaki, 2005; Agarwal et al., 2006, 2010). A rapid and transient increase in *DREB2* expression causes an increase in the regulatory and/or functional proteins which shows the upstream role of *DREB2* in a signaling pathway and improves plant tolerance to

short-time dehydration. Recently, the effect of some PGPRs such as *Arthrobacter protophormiae* (SA3), *Dietzia natronolimnaea* (STR1) and *Bacillus subtilis* (LDR2) on *TaCTR1/TaDREB2* expression (Barnawal et al., 2017) and improvement of wheat tolerance to abiotic stress has been reported but, the effect of *A. brasilense* on *DREB2* expression is not well studied in inoculated wheat (*T. aestivum* L.) cultivars under dehydration condition.

## Materials and methods

### Primary experiment

A primary experiment (unpublished data) was conducted to choose the most tolerant and sensitive cultivars to drought as well as drought tolerant and sensitive associated pairs among the combination of six wheat cultivars of common wheat (*T. aestivum* L.) named Roshan, Sardari, Tabasi, Shahpasand, Omid and Shoaleh inoculated with two strains of *A. brasilense* called Sp7 (as standard strain) and Sp245 (reported to produce significant ABA). The seeds were prepared from Agriculture and Natural Resources Research Center, Isfahan, Iran. The strains of *A. brasilense* strains were obtained from Research Association of AMP Research Unit Microbe-Plant Interactions, Germany.

The growth and biochemical parameters of six wheat cultivars as well as their combinations with two strains of *A. brasilense* (Sp7 and Sp245) including roots and shoots dry weight (RDW, SDW), average roots length (ARL), number of main roots (NMR) and the number of root branches (NRB), maximum shoot length (MSL) as well as total protein of roots and shoots (RTP, STP), PAL enzyme activity in roots and shoots (RPAL, SPAL) and activity of TAL enzyme in roots and shoots (RTAL, STAL) were measured. Adopted method of Bradford (1976) was used for protein assay and the method of Beaudoin-Egan and Thorpe (1985) was used for TAL and PAL enzymes activity analysis. The experiment was conducted under drought condition (40% of field capacity) and compared to 80% moisture content (data were not shown). Then, the data was ranked for the best and the worst responses to drought (*Appendix 1*). Afterward, according to the measured parameters, the two cultivars of Roshan and Shahpasand were selected as drought tolerant and sensitive cultivars, respectively. Simultaneously, the associated pairs of cultivar-strain including Roshan-Sp245, Roshan-Sp7, Shahpasand-Sp7 and, Shahpasand-Sp245 were selected as tolerant, moderately tolerant, sensitive and moderately sensitive to drought within all possible combinations of wheat cultivars and *A. brasilense* strains (*Appendix 1*). Then, in the second experiment, the selected cultivars and associated pairs were evaluated for *DREB2* gene expression under optimum ( $\psi_w = 0$  MPa) and dehydration ( $\psi_w = -90$  MPa) conditions.

### Second experiment

*A. brasilense* strains were cultured on solid NFB (Nitrogen free basal) medium and incubated at 32°C. Then the bacteria were cultured in liquid NFB medium enriched with ammonium chloride (0.1% w/v, 1 g L<sup>-1</sup>) and incubated at 30°C for 48 h. Bacterial concentration of 10<sup>8</sup> CFU mL<sup>-1</sup> was adjusted by reading optical density at 600 nm using McFarland table (Krieg and Dobereiner, 1984; McFarland, 1907).

Seeds were sterilized using the method adopted by Sweet and Bolton (1979). Then the sterilized seeds were incubated in sterile distilled water at room temperature for 3 h. Wheat seeds were shaken in high phosphate Nfb (HPNfb) liquid medium enriched with

0.1% ammonium chloride contain  $10^8$  CFU mL<sup>-1</sup> of *A. brasilense* stain and shake in 100 rpm for 3 h (Shaker Model INFORS AG, BOTTMINGEN) (Okon et al., 1977). Under sterile laminar airflow, the seeds were left to reduce the seeds moisture down to almost 14%.

Sterilized wheat seeds (Shahpasand as sensitive and Roshan as tolerant cultivar) were planted on sterile water agar (8% w/v) and kept in dark at room temperature for 48 h for germination. The uniform small seedlings were transferred into test tube (3 × 25 cm) containing 40 ml of autoclaved MS medium (30 g sucrose, 7 g agar, pH ~ 6.5) (Murashige and Skoog, 1962) and then located in growth chamber at 22 ± 2°C, 60% RH, and photoperiod of 16 L/8 D (210 photons m<sup>-2</sup> s<sup>-1</sup>) for 3 more days. Half of the seedlings (5-day-old) were inoculated with specific *A. brasilense* strains.

To inoculate seedling, Nfb liquid medium enriched with 0.1% ammonium chloride containing  $10^7$  CFU ml<sup>-1</sup> concentration of *A. brasilense* Sp245 and Sp7 was prepared. Then under sterile laminar airflow, 1 mL of Nfb liquid medium containing Sp245 or Sp7 was added separately into each test tube, containing three 5-day-old seedlings and kept for 2 more days. After 2 days, seedling's roots were checked for the number of bacteria. To do so, the samples of seedling's roots were stained with Iodonitrotetrazolium chloride (INT) dye (0.25% w/v aqueous solution) and then many cross-sections of root were provided manually (Sriskandarajah et al., 1993). Afterward, the presence of the bacteria within root cross-sections was checked using a light microscope. Meanwhile, the number of bacteria colonies (CFU mL<sup>-1</sup>) within the roots was counted. To do so, the root samples were placed in 95% (v/v) ethanol for 3 sec, then in 0.01% (w/v) HgCl<sub>2</sub> for 15 sec and washed several times with sterile water for surface-sterilization (Sriskandarajah et al., 1993). Then, 1 g of root samples were grounded using 5 mL of 1X phosphate buffered saline (PBS) (pH ~ 7.4) at the sterile condition and serially diluted ( $10^{-1}$  up to  $10^{-9}$ ) (Stets et al., 2015). 5 µl of the final dilution ( $10^{-9}$ ) were cultured on solid medium (8% w/v agar) of HPNfb enriched with ammonium chloride (0.1% w/w) and incubated at 32°C for 72 h. At last, the number of colonies formed on the plates was counted and considered as bacterial population in 1 g of tissue fresh weight. The number of bacteria in root was almost  $10^8$  CFU mL<sup>-1</sup> for Sp245 and  $10^7$  CFU mL<sup>-1</sup> for Sp7.

One series of 7-day-old seedlings (including non-inoculated and inoculated with *A. brasilense* strains) were transferred into sterile test tubes (3 × 25 cm) which had been saturated with water vapors (RH ≈ 100%) at 25°C (Thomson, 1872). According to Kelvin equation and after equilibrium of water vapor pressure, this was equal to maximum water potential ( $\psi_w = 0$  MPa) and was considered as optimum water condition (Butt et al., 2006). The second series of 7-day-old seedlings were transferred into sterile test tubes with RH equal to 50% ( $\psi_w = -90$  MPa) at 25°C as dehydration treatment. After implementation and equilibrium of treatments, plant samples were collected at 0, 120 and 360 minutes. Samples were separated immediately into shoot and root and frozen in liquid nitrogen to store at -80°C for RNA isolation.

### **Total RNA extraction and quantitative Real-Time PCR (qRT-PCR) analysis**

Total RNA of the shoot and root samples of wheat cultivars were extracted using Irizol kit (Irizol RNA Biotech, Iran) at 4°C. To remove the possible DNA contamination, the RNA samples was treated with RNase-free DNase I (TaKaRa, Japan).

About 1 µg of total RNA was used to synthesize first-strand cDNA primed with oligo (dT) in a 10 µl reaction mix using reverse transcriptase (TaKaRa, Japan) according to the instructions of the manufacturer. QRT-PCR was performed using SYBER-Green dye (TaKaRa, Japan) on real-time PCR machine (Bio-Rad, USA) in triplicate. The *ARF* (ADP-ribosylation factor) gene (AB050957) was used as the reference gene. The primers (*ARF*, F: 5'-CTGACGCCGAGGATATCCA-3'; R: 5'-GCCTTGACCATAACCAGTTCCA-3') used for Real-Time PCR analysis were designed using AlleleID, version 7. The amount of transcript accumulated for the *DREB2* as the target gene (AY781345) (*DREB2*, F: 5'-AAGAAAACAGGCGACAAGAT-3'; and R: 5'-GTTGTTGGTTCACCTTCTTTC-3') (Egawa et al., 2006) normalized to *ARF* gene as internal control. The value of *DREB2* transcript level was reported relative to the *ARF* gene ( $-\Delta C_T$ ) which is the negative difference between the threshold cycle ( $C_T$ ) of the target gene and  $C_T$  of the reference gene. Then, the relative expression of *DREB2* in treated conditions (inoculated and/or dehydrated) was calculated using the  $2^{-\Delta\Delta C_T}$  method relative to control condition (non-inoculation and optimum water condition) (Livak and Schmittgen, 2001). The real-time reaction efficiencies of the target and reference genes ( $E = 0.93$  and  $0.96$ , respectively) were assumed to be equal, nearly 1. The PCR cycling conditions including an optimum concentration of primers and cDNA as well as optimum temperature for *ARF* and *DREB2* genes were set up. So, real-time PCR program was: incubation at 95°C for 10 min, 95°C for 10 s, 58°C for 25 s for 40 cycles, 72°C for 30 s and 72°C for 7 min.

### **Statistical analysis**

The primary experiment was conducted in a split plot layout with 3 replicates. The main plot was wheat cultivars, sub-plot was bacterial strains and sub-sub-plot was water condition. The second experiment also carried out as split-plot with 3 replicates. The main plot was sensitivity to drought, inoculation as sub-plot and water condition as sub-sub-plot.

The ANOVA were obtained using SPSS and the mean values were compared using Duncan's multiple range tests at 95% confidence interval.

### **Results**

According to the primary experiment, the associated pairs (wheat cultivars-*A. brasilense* strains) including tolerant (Roshan-Sp245), moderately tolerant (Roshan-Sp7), sensitive (Shahpasand-Sp7) and moderately sensitive (Shahpasand-Sp245) to drought were selected (*Appendix 1*). Physiological responses in drought condition are presented in *Tables 1 and 2*.

### **Physiological responses of sensitive and tolerant associated pairs to drought**

Cross-inoculation of Roshan and Shahpasand cultivars with Sp245 or Sp7 strains formed different associated pairs with different physiological responses in optimum water condition ( $P \leq 0.05$ ) (*Tables 1-4*). In optimum water condition, most physiological parameters increased (except for NMR and SPAL) ( $P \leq 0.05$ ) when inoculated tolerant cultivar (Roshan- Sp245) compared to non-inoculated one (Roshan). However, in Roshan-Sp7, ARL, RDW and, RTP were the only parameters positively affected by inoculation in optimum water condition (*Tables 3 and 4*). Although,

inoculation of the sensitive cultivar (Shahpasand) with Sp7 did not significantly affect any of the physiological parameters at optimum water condition in Shahpasand-Sp245, NRB, SDW, RTP and, RPAL increased due to inoculation ( $P \leq 0.05$ ) (Tables 1-4). It seems that Sp245 has more homology or compatibility potential to associate with different wheat cultivars than Sp7.

**Table 1.** Average values of growth parameters including maximum shoot length (MSL), average root length (ARL), number of main roots (NMR), number of root branches (NRB) and shoot and root dry weight (SDW and RDW, respectively), in non-inoculated cultivars (Roshan and Shahpasand as tolerant and sensitive to drought, respectively) as well as different pairs of wheat cultivars and *A. brasilense* strains (Sp245 and Sp7) including Roshan-Sp245 (tolerant), Roshan-Sp7 (moderately tolerant), Shahpasand-Sp7 (sensitive), Shahpasand-Sp245 (moderately sensitive) under drought (40% FC) and optimum water conditions (80% FC). Data represent the mean values  $\pm$  SE ( $n = 3$ ). Different letters in the mean values indicated the difference between the mean values (Duncan's multiple range tests,  $P \leq 0.05$ ). MSL and ARL are in cm plant<sup>-1</sup>; SDW and RDW are in mg plant<sup>-1</sup>. FC, stand for moisture content at field capacity

FC	Cultivar, Strain	MSL	ARL	NMR	NRB	SDW	RDW
80%	Roshan cv.	25.3 $\pm$ 0.2 <sup>b</sup>	5.7 $\pm$ 0.2 <sup>c</sup>	5.0 $\pm$ 0.01 <sup>a</sup>	0.7 $\pm$ 0.6 <sup>c</sup>	25.0 $\pm$ 0.01 <sup>b</sup>	9.0 $\pm$ 0.01 <sup>c</sup>
	Roshan-Sp245	31.2 $\pm$ 0.8 <sup>a</sup>	10.6 $\pm$ 0.1 <sup>a</sup>	5.3 $\pm$ 0.6 <sup>a</sup>	2.3 $\pm$ 0.6 <sup>b</sup>	29.3 $\pm$ 0.6 <sup>a</sup>	17.3 $\pm$ 0.6 <sup>a</sup>
	Roshan-Sp7	26.3 $\pm$ 0.3 <sup>b</sup>	7.4 $\pm$ 0.2 <sup>b</sup>	5.0 $\pm$ 0.01 <sup>a</sup>	1.3 $\pm$ 0.6 <sup>c</sup>	26.0 $\pm$ 1.0 <sup>b</sup>	12.7 $\pm$ 0.6 <sup>b</sup>
	Shahpasand cv.	17.8 $\pm$ 0.3 <sup>c</sup>	7.1 $\pm$ 0.1 <sup>b</sup>	4.0 $\pm$ 1.0 <sup>b</sup>	0.0 $\pm$ 0.01 <sup>d</sup>	20.7 $\pm$ 0.6 <sup>c</sup>	7.0 $\pm$ 0.01 <sup>c</sup>
	Shahpasand-Sp245	19.7 $\pm$ 0.6 <sup>c</sup>	7.6 $\pm$ 0.1 <sup>b</sup>	4.0 $\pm$ 0.01 <sup>b</sup>	1.0 $\pm$ 0.01 <sup>c</sup>	24.0 $\pm$ 1.0 <sup>b</sup>	7.0 $\pm$ 0.01 <sup>c</sup>
	Shahpasand-Sp7	18.8 $\pm$ 0.3 <sup>c</sup>	7.4 $\pm$ 0.1 <sup>b</sup>	3.7 $\pm$ 0.6 <sup>b</sup>	0.7 $\pm$ 0.6 <sup>c</sup>	22.0 $\pm$ 0.01 <sup>c</sup>	7.0 $\pm$ 0.01 <sup>c</sup>
40%	Roshan cv.	24.0 $\pm$ 0.2 <sup>b</sup>	5.4 $\pm$ 0.1 <sup>c</sup>	5.0 $\pm$ 0.01 <sup>a</sup>	2.0 $\pm$ 0.01 <sup>b</sup>	24.0 $\pm$ 0.01 <sup>b</sup>	8.0 $\pm$ 0.01 <sup>c</sup>
	Roshan-Sp245	26.4 $\pm$ 0.5 <sup>b</sup>	7.4 $\pm$ 0.4 <sup>b</sup>	5.0 $\pm$ 0.01 <sup>a</sup>	3.7 $\pm$ 0.6 <sup>a</sup>	27.7 $\pm$ 0.6 <sup>a</sup>	12.0 $\pm$ 0.01 <sup>b</sup>
	Roshan-Sp7	24.3 $\pm$ 0.2 <sup>b</sup>	6.2 $\pm$ 0.1 <sup>b</sup>	5.0 $\pm$ 0.01 <sup>a</sup>	2.3 $\pm$ 0.6 <sup>b</sup>	25.0 $\pm$ 0.01 <sup>b</sup>	9.0 $\pm$ 0.01 <sup>c</sup>
	Shahpasand cv.	13.8 $\pm$ 0.5 <sup>d</sup>	5.4 $\pm$ 0.1 <sup>c</sup>	4.3 $\pm$ 0.6 <sup>b</sup>	2.0 $\pm$ 0.01 <sup>b</sup>	16.0 $\pm$ 1.0 <sup>d</sup>	5.0 $\pm$ 0.01 <sup>d</sup>
	Shahpasand-Sp245	13.5 $\pm$ 0.5 <sup>d</sup>	5.4 $\pm$ 0.1 <sup>c</sup>	4.7 $\pm$ 0.6 <sup>ab</sup>	1.3 $\pm$ 0.6 <sup>c</sup>	16.0 $\pm$ 1.0 <sup>d</sup>	5.0 $\pm$ 0.01 <sup>d</sup>
	Shahpasand-Sp7	9.4 $\pm$ 0.5 <sup>e</sup>	3.8 $\pm$ 0.1 <sup>d</sup>	3.0 $\pm$ 0.01 <sup>c</sup>	1.7 $\pm$ 0.6 <sup>b</sup>	11.0 $\pm$ 1.7 <sup>e</sup>	2.0 $\pm$ 0.01 <sup>e</sup>

**Table 2.** Average values of total protein in shoots (STP) and roots (RTP), activity of PAL enzyme in shoots (SPAL) and roots (RPAL), activity of TAL enzyme in shoot (STAL) and root (RTAL) in non-inoculated cultivars (Roshan and Shahpasand as tolerant and sensitive to drought, respectively) as well as different pairs of wheat cultivars and *A. brasilense* strains (Sp245 and Sp7) including Roshan-Sp245 (tolerant), Roshan-Sp7 (moderately tolerant), Shahpasand-Sp7 (sensitive), Shahpasand-Sp245 (moderately sensitive) under drought (40% FC) and optimum water conditions (80% FC). Data represent the mean values  $\pm$  SE ( $n = 3$ ). Different letters in the mean values indicated the difference between the means (Duncan's multiple range tests,  $P \leq 0.05$ ). STP and RTP are in mg gFW<sup>-1</sup>; SPAL, RPAL, STAL and RTAL are in U mg protein<sup>-1</sup>. FC, stand for moisture content at field capacity

FC	Cultivar, Strain	STP	RTP	SPAL	RPAL	STAL	RTAL
80%	Roshan cv.	44.0 $\pm$ 0.5 <sup>b</sup>	37.4 $\pm$ 1.1 <sup>c</sup>	17.4 $\pm$ 0.01 <sup>c</sup>	11.7 $\pm$ 0.2 <sup>b</sup>	20.7 $\pm$ 0.1 <sup>c</sup>	15.3 $\pm$ 0.01 <sup>d</sup>
	Roshan-Sp245	86.2 $\pm$ 0.5 <sup>a</sup>	79.3 $\pm$ 1.0 <sup>a</sup>	20.6 $\pm$ 0.1 <sup>b</sup>	14.5 $\pm$ 0.3 <sup>a</sup>	23.6 $\pm$ 0.1 <sup>b</sup>	19.3 $\pm$ 0.6 <sup>c</sup>
	Roshan-Sp7	45.0 $\pm$ 0.2 <sup>b</sup>	47.3 $\pm$ 0.7 <sup>bc</sup>	18.0 $\pm$ 0.1 <sup>c</sup>	12.4 $\pm$ 0.1 <sup>b</sup>	21.9 $\pm$ 0.1 <sup>c</sup>	16.0 $\pm$ 0.4 <sup>d</sup>
	Shahpasand cv.	32.7 $\pm$ 0.3 <sup>c</sup>	26.0 $\pm$ 1.3 <sup>d</sup>	10.8 $\pm$ 0.1 <sup>d</sup>	9.0 $\pm$ 0.01 <sup>c</sup>	19.4 $\pm$ 0.1 <sup>c</sup>	12.1 $\pm$ 0.3 <sup>e</sup>
	Shahpasand-Sp245	37.6 $\pm$ 0.5 <sup>c</sup>	35.0 $\pm$ 1.5 <sup>c</sup>	12.3 $\pm$ 0.2 <sup>d</sup>	11.5 $\pm$ 0.3 <sup>b</sup>	21.0 $\pm$ 0.1 <sup>c</sup>	13.3 $\pm$ 0.3 <sup>e</sup>
	Shahpasand-Sp7	36.1 $\pm$ 1.2 <sup>c</sup>	29.2 $\pm$ 1.2 <sup>d</sup>	11.1 $\pm$ 0.3 <sup>d</sup>	9.9 $\pm$ 0.1 <sup>c</sup>	19.7 $\pm$ 0.1 <sup>c</sup>	12.5 $\pm$ 0.3 <sup>e</sup>
40%	Roshan cv.	50.9 $\pm$ 0.9 <sup>b</sup>	56.1 $\pm$ 4.6 <sup>b</sup>	23.6 $\pm$ 0.6 <sup>a</sup>	12.9 $\pm$ 0.1 <sup>b</sup>	24.8 $\pm$ 0.2 <sup>b</sup>	18.3 $\pm$ 0.6 <sup>c</sup>
	Roshan-Sp245	55.7 $\pm$ 0.7 <sup>b</sup>	79.3 $\pm$ 0.0 <sup>a</sup>	26.2 $\pm$ 0.1 <sup>a</sup>	15.4 $\pm$ 0.01 <sup>a</sup>	30.3 $\pm$ 0.1 <sup>a</sup>	25.0 $\pm$ 0.7 <sup>a</sup>
	Roshan-Sp7	52.0 $\pm$ 0.4 <sup>b</sup>	70.3 $\pm$ 1.8 <sup>a</sup>	24.2 $\pm$ 0.1 <sup>a</sup>	13.5 $\pm$ 0.3 <sup>b</sup>	26.0 $\pm$ 0.1 <sup>b</sup>	22.3 $\pm$ 1.2 <sup>b</sup>
	Shahpasand cv.	30.3 $\pm$ 0.3 <sup>c</sup>	21.5 $\pm$ 1.1 <sup>d</sup>	9.7 $\pm$ 0.1 <sup>d</sup>	8.1 $\pm$ 0.1 <sup>ce</sup>	17.9 $\pm$ 0.1 <sup>c</sup>	10.2 $\pm$ 0.3 <sup>e</sup>
	Shahpasand-Sp245	32.2 $\pm$ 0.7 <sup>c</sup>	21.2 $\pm$ 0.2 <sup>d</sup>	8.6 $\pm$ 0.1 <sup>d</sup>	7.2 $\pm$ 0.3 <sup>e</sup>	16.6 $\pm$ 0.1 <sup>c</sup>	9.20 $\pm$ 0.3 <sup>ef</sup>
	Shahpasand-Sp7	32.3 $\pm$ 0.3 <sup>c</sup>	19.4 $\pm$ 0.9 <sup>d</sup>	8.3 $\pm$ 0.01 <sup>d</sup>	7.0 $\pm$ 0.2 <sup>e</sup>	15.2 $\pm$ 0.1 <sup>d</sup>	6.5 $\pm$ 0.6 <sup>f</sup>

**Table 3.** Based on the values of Table 1, the percent of changes in each parameter of different associated pairs due to inoculation and/or drought conditions. Optimum water condition defined as 80% FC and drought as 40% FC (moisture content at field capacity). Abbreviation: maximum shoot length (MSL), average roots length (ARL), number of main roots (NMR), number of roots branches (NRB) and shoots and roots dry weight (SDW and RDW, respectively)

Growth parameters	Sensitivity of associated pairs to drought	At 80% FC Inoculated versus Non-inoculated	For non-inoculated 40% versus 80%	For Inoculated 40% versus 80%	Inoculated 40% versus Non-inoculated 80%
		Inoculation effect	Drought effect	Drought effect under inoculation	Dual effect
MSL	Tolerant	23	0	-15	0
	Moderately tolerant	0	-	0	0
	Sensitive	0	-25	-50	-47
	Moderately sensitive	0	-	-31	-24
ARL	Tolerant	85	0	-30	29
	Moderately tolerant	30	-	0	9
	Sensitive	0	-25	-50	-46
	Moderate sensitive	0	-	-29	-24
NMR	Tolerant	0	0	0	0
	Moderately tolerant	0	-	0	0
	Sensitive	0	0	-18	-25
	Moderately sensitive	0	-	0	0
NRB	Tolerant	250	198	61	448
	Moderately tolerant	0	-	77	229
	Sensitive	0	2*	143	1.67*
	Moderately sensitive	1*	-	0	1.3*
SDW	Tolerant	16	0	0	12
	Moderately tolerant	0	-	0	0
	Sensitive	0	-25	-50	-47
	Moderately sensitive	16	-	-33	-23
RDW	Tolerant	89	0	-31	33
	Moderately tolerant	41	-	-29	0
	Sensitive	0	-25	-71	-71
	Moderately sensitive	0	-	-28	-28

\*: Indication increased NRB from 0

The selected cultivars or associated pairs showed different physiological responses to drought ( $P \leq 0.05$ ) (Tables 1-4). At drought condition, MSL, ARL and, RDW in Roshan cv. (drought tolerant) did not decrease as compared to optimum water condition, but these parameters were decreased in Shahpasand cv. (Tables 1 and 3). In drought condition, RDW decreased in all pairs with the highest in Shahpasand-Sp7 and the lowest in Roshan-Sp245. SDW of Roshan cv. in non-inoculated and inoculated status did not change in drought condition but SDW of Shahpasand cv. and inoculated Shahpasand cv. with any of two strains were decreased. The drought condition had a negative effect on NMR of Shahpasand with Sp7. However, NRB increased in all cultivars and pairs except for Shahpasand-Sp245 under drought condition ( $P \leq 0.05$ ) (Tables 1-4).

STAL and RTAL were increased in inoculated Roshan cv. with Sp245 or Sp7 but they decreased in Shahpasand cv. with both strains. In contrast, SPAL in Roshan cv. and Roshan inoculated with Sp245 or Sp7 were increased in drought condition but they did

not change in Shahpasand cv. or Shahpasand with Sp7 or Sp245. STP did not change due to drought in all cultivars and pairs except for Roshan-Sp245. Meanwhile, RTP did not change in Roshan-Sp245 and Shahpasand-Sp7 but it was increased in Roshan-Sp7 and decreased in Shahpasand-Sp245 under drought condition (Tables 2 and 4).

**Table 4.** Based on the values of Table 2, the percent of changes in each parameter of different associated pairs due to inoculation and/or drought conditions. Optimum water condition defined as 80% FC and drought as 40% FC (moisture content at field capacity). Abbreviation: total protein of roots (RTP) and shoots (STP), the activity of PAL enzyme in shoots (SPAL) and roots (RPAL), the activity of TAL enzyme in shoots (STAL) and roots (RTAL)

Growth parameters	Sensitivity of associated pairs to drought	At 80% FC Inoculated versus Non-inoculated	For non-inoculated	For Inoculated	Inoculated 40% versus Non-inoculated 80%
			40% versus 80%	40% versus 80%	
		Inoculation effect	Drought effect	Drought effect under inoculation	Dual effect
STP	Tolerant	96	0	-35	0
	Moderately tolerant	0	-	0	0
	Sensitive	0	0	0	0
	Moderately sensitive	0	-	0	0
RTP	Tolerant	110	50	0	112
	Moderately tolerant	26	-	49	88
	Sensitive	0	0	0	0
	Moderately sensitive	35	-	-39	0
SPAL	Tolerant	0	36	27	50
	Moderately tolerant	0	-	34	39
	Sensitive	0	0	0	0
	Moderately sensitive	0	-	0	0
RPAL	Tolerant	24	0	0	31
	Moderately tolerant	0	-	0	0
	Sensitive	0	0	-29	-29
	Moderately sensitive	28	-	-37	-20
STAL	Tolerant	14	20	29	47
	Moderately tolerant	0	-	19	26
	Sensitive	0	0	-23	-22
	Moderately sensitive	0	-	-37	0
RTAL	Tolerant	27	20	29	37
	Moderately tolerant	0	-	39	46
	Sensitive	0	0	-48	-47
	Moderately sensitive	0	-	-31	-24

The dual effects of inoculation and drought were evaluated by comparing an inoculated cultivar which was grown under drought condition with the respective non-inoculated cultivar at optimum water condition (named control). When inoculated Roshan cv. with Sp245 or Sp7 under drought condition were compared with control, ARL, NRB, RTP, SPAL, RTAL and, STAL were higher in inoculated one. In contrast, RDW, SDW, ARL, MSL, RPAL, RTAL in Shahpasand cv. with Sp7 or Sp245 under drought condition were lower than control. Simultaneously, MSL, NMR and STP in inoculated Roshan cv. and RTP, STP and, SPAL of inoculated Shahpasand cv. did not show any changes. However, NRB increased in inoculated Shahpasand cv. as compared to control (Tables 1-4).



In brief, the most beneficial changes in the physiological parameters were observed in the tolerant pair (Roshan-Sp245) followed by moderately tolerant (Roshan-Sp7), moderately sensitive (Shahpasand- Sp245) and sensitive pairs (Shahpasand-Sp7) at either drought or optimum water conditions (Tables 1-4).

### ***DREB2* expression in selected pairs under different water potentials**

At optimum water potential ( $\psi_w = 0$  MPa), *DREB2* expression in the roots of inoculated seedlings (Tables 5, 6) was less relative to non-inoculated cultivars (ranged from 0.05 up to 0.57- fold for tolerant to sensitive pairs ( $P \leq 0.05$ )). However, their expressions were similar at 0, 120 and 360 min after dehydration imposed (Table 5). *DREB2* in the shoots of all selected pairs were up-regulated as compared to non-inoculated cultivars. *DREB2* up-regulation was more in the shoots of tolerant and moderately tolerant pairs than the sensitive and moderately sensitive pairs ( $P \leq 0.05$ ) (Table 6). Therefore, at optimum water condition, *DREB2* down-regulated in the roots (Table 5) but up-regulated in the shoots of tolerant and sensitive pairs relative to non-inoculated cultivars.

**Table 5.** *DREB2* expression (fold) in different times (0, 120 and 360 min) after dehydration ( $\psi_w = -90$  MPa) imposed in the roots of different associated pairs of wheat cultivars and *A. brasilense* strains (Sp245 and Sp7) as responded to drought including: tolerant (Roshan-Sp245), moderately tolerant (Roshan-Sp7), sensitive (Shahpasand-Sp7), moderately sensitive (Shahpasand-Sp245). The result was compared to optimum water condition ( $\psi_w = 0$  MPa). Data represent the mean values  $\pm$  SE ( $n = 3$ ). Different letters in the mean values indicated the difference between the means (Duncan's multiple range tests,  $P \leq 0.05$ )

Time (min)	$\Psi_w$ (MPa)	Sensitivity of associated pairs to drought	Non-inoculated	Inoculated	Inoculated versus Non-inoculated
			Dehydration versus optimum	Dehydration versus optimum	
0	0 (optimum)	Tolerant	-	-	$0.06 \pm 0.02^c$
		Moderately tolerant	-	-	$0.05 \pm 0.01^c$
		Sensitive	-	-	$0.15 \pm 0.04^c$
		Moderately sensitive	-	-	$0.15 \pm 0.02^c$
120	0	Tolerant	-	-	$0.07 \pm 0.02^c$
		Moderately tolerant	-	-	$0.06 \pm 0.02^c$
		Sensitive	-	-	$0.14 \pm 0.02^c$
		Moderately sensitive	-	-	$0.16 \pm 0.01^c$
	-90 (dehydrated)	Tolerant	$2.65 \pm 0.70^a$	$777.8 \pm 100.76^a$	$19.20 \pm 4.83^a$
		moderately tolerant	-	$710.8 \pm 84.44^a$	$17.19 \pm 2.12^b$
		Sensitive	$1.60 \pm 0.02^b$	$20.9 \pm 1.16^b$	$1.87 \pm 0.23^c$
		Moderately sensitive	-	$19.1 \pm 2.72^b$	$1.86 \pm 0.42^c$
360	0	Tolerant	-	-	$0.07 \pm 0.01^c$
		Moderately tolerant	-	-	$0.06 \pm 0.02^c$
		Sensitive	-	-	$0.20 \pm 0.52^c$
		moderately sensitive	-	-	$0.20 \pm 0.11^c$
	-90	Tolerant	$0.51 \pm 0.14^c$	$3.6 \pm 0.68^b$	$0.51 \pm 0.06^c$
		Moderately tolerant	-	$3.4 \pm 0.43^b$	$0.43 \pm 0.10^c$
		Sensitive	$1.93 \pm 0.48^b$	$5.1 \pm 0.56^b$	$0.51 \pm 0.09^c$
		Moderately sensitive	-	$5.5 \pm 5.06^b$	$0.57 \pm 0.02^c$

Dehydration had a significant effect on *DREB2* expression of roots and shoots in inoculated and non-inoculated cultivars ( $P \leq 0.05$ ) (Tables 5, 6). The result showed that in the tolerant cultivar (Roshan cv.) at 120 min after dehydration, *DREB2* ( $2^{-\Delta\Delta C_T}$ ) up-regulated 2.7 and 3.4- fold in the roots and shoots, respectively as compared to optimum

water condition. In the sensitive cultivar (Shahpasand cv.), *DREB2* did not change in the roots or shoots due to dehydration ( $P \leq 0.05$ ) (Table 5). At 360 min, *DREB2* expression acts vice versa under dehydration, means, down-regulated in the roots of Roshan (0.5-fold) and up-regulated (2.6- fold) in the shoots, but in Shahpasand, *DREB2* up-regulated in the roots (1.9- fold) and down-regulated in the shoots (0.6- fold).

The change in *DREB2* expression due to dehydration in non-inoculated cultivars, and selected association pairs (the inoculated cultivars) had different sensitivity to drought (Table 5). For example, at 120 min after dehydration, *DREB2* up-regulated in the roots of tolerant, moderately tolerant, sensitive and moderately sensitive pairs (778, 711, 21 and 19- fold, respectively) while, *DREB2* up-regulated (1-fold) in the shoots of tolerant and moderately tolerant pairs and down-regulated (almost 0.4- fold) in the sensitive and moderately sensitive pairs (Tables 5, 6). However, at 360 min, *DREB2* up-regulation in the roots of all pairs were so less (ranged from 3.4 to 5.5- fold). In contrast, *DREB2* down-regulated due to dehydration in the shoots of the tolerant, moderately tolerant (0.2- fold), sensitive and moderately sensitive pairs (0.4- fold) (Tables 5, 6).

**Table 6.** *DREB2* expression (fold) in different times (0, 120 and 360 min) after dehydration ( $\psi_w = -90$  MPa) imposed in the shoots of different associated pairs of wheat cultivars and *A. brasilense* strains (Sp245 and Sp7) as responded to drought including: tolerant (Roshan-Sp245), moderately tolerant (Roshan-Sp7), sensitive (Shahpasand-Sp7), moderately sensitive (Shahpasand-Sp245). The result was compared to optimum water condition ( $\psi_w = 0$  MPa). Data represent the mean values  $\pm$  SE ( $n = 3$ ). Different letters in the mean values indicated the difference between the means (Duncan's multiple range tests,  $P \leq 0.05$ )

Time (min)	$\Psi_w$ (MPa)	Sensitivity of associated pairs to drought	Non-inoculated	Inoculated	Inoculated versus Non-inoculated
			Dehydration versus optimum	Dehydration versus optimum	
0	0 (control)	Tolerant	-	-	$3.43 \pm 0.47^a$
		Moderately tolerant	-	-	$3.27 \pm 0.70^a$
		Sensitive	-	-	$1.39 \pm 0.32^b$
		Moderately sensitive	-	-	$1.40 \pm 0.12^b$
120	0	Tolerant	-	-	$3.35 \pm 0.64^a$
		Moderately tolerant	-	-	$3.32 \pm 0.19^a$
		Sensitive	-	-	$1.27 \pm 0.01^b$
		Moderately sensitive	-	-	$1.28 \pm 0.03^b$
	-90 (dehydrated)	Tolerant	$3.35 \pm 0.33^a$	$1.22 \pm 0.29^a$	$1.22 \pm 0.32^b$
		moderately tolerant	-	$1.12 \pm 0.17^a$	$1.11 \pm 0.19^b$
		Sensitive	$1.02 \pm 0.31^b$	$0.38 \pm 0.14^b$	$0.48 \pm 0.18^d$
		Moderately sensitive	-	$0.40 \pm 0.09^b$	$0.50 \pm 0.06^d$
360	0	Tolerant	-	-	$3.29 \pm 1.21^a$
		Moderately tolerant	-	-	$3.08 \pm 0.85^a$
		Sensitive	-	-	$1.17 \pm 0.02^b$
		moderately sensitive	-	-	$1.18 \pm 0.34^b$
	-90	Tolerant	$2.64 \pm 0.47^a$	$0.16 \pm 0.04^c$	$0.20 \pm 0.05^e$
		Moderately tolerant	-	$0.16 \pm 0.08^c$	$0.19 \pm 0.05^e$
		Sensitive	$0.59 \pm 0.09^b$	$0.43 \pm 0.14^b$	$0.85 \pm 0.18^c$
		Moderately sensitive	-	$0.45 \pm 0.09^b$	$0.90 \pm 0.26^c$

To evaluate dual effects of inoculation and dehydration in each pair, the inoculated seedlings which exposed to dehydration condition were compared to the respective non-inoculated seedlings grown at optimum water condition. The result showed that at 120 min after dehydration, transcripts level of *DREB2* ( $-\Delta CT$ ) in the tolerant and moderately

tolerant pairs was more than tolerant cultivar grown at optimum water condition (11.4- fold for roots and 2.3- fold for shoots). Meanwhile, in dehydrated roots of sensitive and moderately sensitive pairs, transcripts level of *DREB2* was more than non-inoculated and non-dehydrated roots (1.5- fold) but in their shoot *DREB2* was less than non-inoculated ones at optimum water potential (2.6 and 2.4- fold, respectively) ( $P \leq 0.05$ ) ( $t = 120$ ).

In the last measurement after dehydration (360 min), the transcripts level of *DREB2* in the tolerant and moderately tolerant pairs decreased (almost 5 and 6-fold for roots and almost 1-fold for shoot, respectively) as compared to non-inoculated cultivars grown under optimum water potential while in the sensitive and moderately sensitive pairs, there was no change in the roots but it was lower in the shoots (3.7 and 3.4-fold, respectively) ( $P \leq 0.05$ ).

## Discussion

The primary experiment was necessary to find out the most sensitive and tolerant cultivars to drought among the usual cultivated wheat cultivars (Roshan, Sardari, Tabasi, Shahpasand, Omid and Shoaleh). Then finding the compatible (or homologous) and incompatible (heterogonous) pairs among association pair of these six cultivars with two strains of *A. brasilense* (Sp7 and Sp245) under drought condition (Baldani et al., 1983; Jain and Patriquin, 1984; Amooaghaie et al., 2004). This is important due to the complexity of interaction between bacteria and host plant which can affect the successful infection, the establishment of bacterial in host plant and the following appropriate physiological responses (Jain and Patriquin, 1984; Sumner, 1990; Baldani et al., 1986, 1987). According to the physiological parameters such as growth and defensive parameters as well as total protein content, the two cultivars of Roshan and Shahpasand were determined as the most tolerant and sensitive cultivars to drought, respectively (Baldani et al., 1983; Estabrook and Sengupta-Gopalan, 1991; Vasse et al., 1993; Santos et al., 2001). Simultaneously, the best partner of wheat cv. and *A. brasilense* strains as tolerant pair to drought was Roshan-Sp245 (homologous) and the worst as the most sensitive pair was Shahpasand-Sp7 (heterologous). Also, other pairs from cross-inoculation of Roshan and Shahpasand cultivars with Sp245 and Sp7 strains indicated that Roshan-Sp7 as moderately tolerant and Shahpasand-Sp245 as moderately sensitive pairs.

### *Physiological responses of the selected pairs to drought*

Cross-inoculation of Roshan and Shahpasand cultivars (as tolerant and sensitive to drought, respectively) with *A. brasilense* Sp245 and Sp7 strains make different associated pairs with different physiological responses in drought and optimum water conditions. Inoculation of Roshan cv. with *A. brasilense* Sp245 improved the growth of wheat seedlings except for NMR and SPAL. Similar results have been reported that Sp245 caused higher shoot length, root length and the number of roots (Baldani et al., 1987; Hamdia et al., 2004; Akbari et al., 2007; Cohen et al., 2008; Spaepen et al., 2009). Also, in the moderately tolerant (Roshan-Sp7) pair, ARL, RDW and RTP increased due to inoculation at optimum water condition (*Tables 3 and 4*). The positive effect of *A. brasilense* to promote wheat growth could be partly related to the production of more roots and consequently uptake more water and nutrient from the soil (Casimiro et al., 2003; Correa-Aragunde et al., 2004). However, inoculation of Shahpasand cv. (sensitive

to drought) with Sp7 (sensitive association pair) did not show any positive effect on the growth parameters but in Shahpasand-Sp245 (moderately sensitive), NRB, SDW, RTP and, RPAL increased due to inoculation. Regarding to previous works, it can be concluded that any strain of bacteria should be homologous to specific wheat cultivar to expect stimulation effect on growth parameter (Baldani et al., 1983; Amooaghaie et al., 2004). In this experiment, not only Sp245 was in favor of tolerant cultivar, but also did a good job with Shahpasand cultivar. In contrast, Sp7 act in favor of Roshan cv. but has no effect on Shahpasand cultivar.

The effect of any bacterial strain on growth and development of wheat cultivars is neither the same nor ended to the growth parameters because in some cases there are no changes in growth indexes but biochemical parameters were affected. For example, total protein of roots (RTP) did not change in Roshan and Shahpasand with Sp7 at optimum water condition, meanwhile, other parameters behave differently (*Tables 2 and 4*). In contrast, RTP in the tolerant and moderately sensitive pairs of Roshan and Shahpasand with Sp245 were increased due to inoculation. An increase in the protein of plants have been reported in an association of different plant species and bacterial strains ((Vasse et al., 1993; Gamas et al., 1998). For example, different strains of *A. brasilense* have been caused an addition of total protein in wheat and tomatoes (Saubidet et al., 2002; Mangmang et al., 2015).

PAL and TAL activity in the roots and shoots of Roshan cv. inoculated with Sp245 were increased compare to non-inoculated cultivars at optimum water condition but there was no change in other association pairs except for RPAL in the moderately sensitive pair (Shahpasand-Sp245) which increased by 28%. *A. brasilense* Sp245 may enhance the defensive mechanism of wheat cultivars via changing the activities of these enzymes same as other PGPRs, then less reduction in growth parameters was observed in Rhoshan-Sp245 compare to other pairs (Maksimov et al., 2015). There is an evidence that an establishment of a proper relation between plant and bacteria can stimulate the biosynthesis pathway of phenolic compounds as a plant defense mechanism (Vasse et al., 1993). TAL and PAL enzymes are involved in the biosynthesis of phenolic compounds via biosynthesis pathway of phenylpropanoids (Lawton et al., 1983; Nemat Alla et al., 2002). Therefore, increase in the activity of TAL and PAL can cause more phenolic compounds in inoculated plants with a homologous bacterial strain which is an efficient mechanism to deal with the adverse effects of reactive oxygen species during the infection process (Estabrook and Sengupta-Gopalan, 1991). This may be a reason for the less damage to the growth rate in this experiment in inoculated as well as drought conditions.

All growth parameters in non-inoculated tolerant cultivar did not change due to drought condition (except more number of root branches, NRB). However, the most growth parameters in sensitive cultivar were severely decreased. Other reports also show that the reduction in growth parameters in the sensitive cultivars of wheat is more than the tolerant cultivars (Liu et al., 2004). Significant reduction in dry weight and other growth parameters of the sensitive cultivars under drought condition may be due to reduced plant water potential and impairment of mineral absorption and less photosynthesis, protein synthesis and carbohydrate metabolism (Tekle and Alemu, 2016). Among the growth parameters, the number of root branches was increased due to drought in both sensitive and tolerant cultivars with the different rate. Development of an extensive root system in drought condition is an avoidance mechanism to obtain more water from the rhizosphere to protect themselves against drought stress (Liu et al.,

2004). The root and shoot elongation not only was observed in drought condition but also osmotic stress can cause higher root length and also higher activity of PAL and TAL enzymes in wheat seedlings inoculated with *A. brasilense* strains (Nabti et al., 2010; Hamdia et al., 2004).

Drought condition caused the addition of total protein in root in non-inoculated tolerant cultivar but no change in the sensitive one as compare to optimum water condition. Addition of total protein due to drought stress indicated the role of organic osmolytes in osmotic adjustment for adaptation of plant to drought stress (Serraj and Sinclair, 2002).

Dual effect of inoculation and drought condition simultaneously caused mostly higher growth parameters in Roshan with Sp245 (except for NMR and MSL) as well as with Sp7 (except ARL and NRB), while the growth parameters in Shahpasand cv. (except for NRB) were significantly decreased as compared to non-inoculated cultivars at optimum water condition. Similar studies showed that inoculation of wheat with Sp245 strain increased growth parameters more than Sp7 due to the capability of Sp245 to colonize interior parts of wheat root unlike the Sp7 (Michiels et al., 1991). The Sp245 strain could improve drought tolerance of wheat by the production of ABA, a key phytohormone in plant response to drought stress (Michiels et al., 1991; Cohen et al., 2008).

Dual effect of inoculation and drought condition show that Roshan cv. with Sp245 caused an increase in PAL and TAL activities compared to non-inoculated cultivars at drought or optimum water condition. It has been suggested that inoculated of wheat cultivars with Sp245 has a positive effect on the activity of these enzymes either at drought or optimum water condition. Hence, *A. brasilense* Sp245 can enhance the defensive mechanism of wheat at the drought condition (Maksimov et al., 2015). PAL and TAL have a main role in drought tolerance in wheat plants through phenylpropanoids biosynthesis pathway and accumulation of more phenolic compounds in plants (Redmann, 1974; Kaur and Zhawar, 2015).

Therefore, it can be concluded that positive effects of inoculation on biochemical properties of the wheat cultivars are mostly related to homology properties of wheat cultivar and bacterial strain not to the sensitivity of cultivar to drought. This is the reason that Roshan (tolerant) and Shahpasand (sensitive) cultivars with Sp245 showed the better results than Sp7. However, the dual effect of inoculation and drought condition represents the interaction effect on the association of wheat cultivar with the strain of bacteria when compared to separate effect.

### ***DREB2* expression of the selected association pairs in different water potentials**

At optimum water condition ( $\psi_w = 0$  MPa), *DREB2* of inoculated cultivars was down-regulated in the root but up-regulated in the shoot of all selected pairs relative to the respective non-inoculated cultivars. It seems that over-expression of *DREB2* in the shoot of inoculated seedlings may be due to the basic effect of *A. brasilense* strains on the wheat seedling. However, the positive effect in the shoot of tolerant and moderately tolerant pairs is more than sensitive and moderately sensitive pairs. The effect of some other PGPRs such as *Arthrobacter protophormiae* (SA3), *Dietzia natronolimnaea* (STR1) and *Bacillus subtilis* (LDR2) on *TaCTR1/TaDREB2* expression in wheat has been shown the enhancement of *TaDREB2* gene expression encoding for a transcription factor and improving the tolerance of plants to abiotic stress conditions (Barnawal et al., 2017).

*DREB2* expression in non-inoculated cultivars and the cross-inoculated with *A. brasilense* strains was also varied under dehydration condition. *DREB2* up-regulated in root and shoot of non-inoculated tolerant cv. but did not change in the sensitive cv. due to dehydration. *DREB2* up-regulation has been reported in dehydration condition in the tolerant cv. of wheat. For example, Shen and colleagues (2003) reported that drought stress increased *TaDREB2* gene expression in tolerant cultivars of wheat. Meanwhile, transgenic *RD29A-GmDREB* wheat plants had a better tolerance compared to wild type plants which shows the contribution of these two genes in wheat drought tolerance (Shiqing et al., 2005). At 360 minutes after dehydration imposed, *DREB2* down-regulated in the root of Roshan and up-regulated in the shoot. As an early gene, *DREB2* expression is rapid and transient (peaking at 2 or 3 hours) and then their mRNA levels will decrease (Yamaguchi-Shinozaki and Shinozaki, 2005; Agarwal et al., 2006, 2010). *DREB2* down-regulation at longer time of dehydration (360 min) in the tolerant cv. may be due to the transient role of this transcription factor. In contrast, in the sensitive cv., *DREB2* up-regulated at this time in the root and down-regulated in the shoot. It seems that in the sensitive cv. at the shorter time (120 min), dehydration could not make *DREB2* up-regulation unless dehydration duration would be longer. Longer dehydration (360 min) means more loss of water potential and more osmotic signals which were produced by the root cells (Taylor et al., 2000; Stein et al., 2000; Rock, 2000; Liu et al., 2005). Likely, the production of signals for stimulation of *DREB2* expression was enough only for the roots of sensitive cv. but not for its shoots. This is the reason for *DREB2* down-regulation even at 360 min after dehydration applied.

In this experiment, up-regulation of *DREB2* in the tolerant and moderately tolerant pairs and its down-regulation in the sensitive and moderately sensitive pairs at dehydration supported the role of *DREB2* gene as an essential early stress-responsive gene in plant tolerance to dehydration (Yamaguchi-Shinozaki and Shinozaki, 2005; Fujita et al., 2011). The early genes encode regulatory proteins such as transcription factors, signal transduction related proteins, osmolytes biosynthesis (Agarwal et al., 2010). There is evidence for better growth of recombinant *E. coli* cells expressing *SbDREB2A* (obtained from *Salicornia brachiata*) in the culture mediums enriched by NaCl, PEG, and mannitol due to the role of plant's *SbDREB2A* in the regulation of other stress-regulated genes. This function represents certain interactions between the plant's *SbDREB2A* with a transcriptional network of the bacterial cells (Gupta et al., 2010). According to our results, it seems that there is probably proper crosstalk between the transcriptional network of *A. brasilense* Sp245 (the most homologous strain) with Roshan (the most proper cultivar of wheat) which causes highest *DREB2* up-regulation in the root of the tolerant pair. *DREB2* up-regulation in the root of tolerant pair was more than the shoot which might be due to the localization of *A. brasilense* Sp245 in the root of Roshan cv. and supports this crosstalk between the transcriptional network of bacteria and plant. *DREB2* up-regulation in the root (778-fold) of tolerant pair was more than shoot (1.2-fold) but in the tolerant cv. *DREB2* expression was less in the root than the shoot under dehydration condition. This result supports the role of Sp245 to change *DREB2* expression in the root of Roshan cultivar. In the sensitive cv., unlike the tolerant cv., *DREB2* expression did not change.

Dual effects of dehydration and inoculation on *DREB2* expression in the tolerant and sensitive pair were different. This effect was positive in the tolerant and moderately tolerant pairs but it was mostly negative or ineffective in the sensitive and moderately sensitive pairs. Meanwhile, the quantity of dual effect was dependent on dehydration

course. According to our result, it is suggested that inoculation of wheat with *A. brasilense* Sp245 can provide a proper tool to improve tolerance of wheat to dehydration via regulation of *DREB* expression.

## Conclusion

In conclusion, the effect of inoculation on physiological (growth and biochemical) parameters of wheat seedlings in the sensitive and moderately sensitive pairs was less than the tolerant and moderately tolerant pairs at drought or optimum water condition. The dual effects of dehydration and inoculation on *DREB2* expression were positive in the tolerant and moderately tolerant pairs and mostly negative or ineffective in the sensitive and moderately sensitive pairs. However, these effects were depending on the duration of dehydration. It seems that dehydration alone could not cause up-regulation of *DREB2* in non-inoculated wheat cultivars unless the time of stress would be enough meanwhile over-expression of *DREB2* in inoculated wheat cultivars with *A. brasilense* after dehydration may be more due to the effect of the bacteria on wheat seedlings. The authors feel further research is needed to find out the details of the effect of *A. brasilense* on *DREB2* gene expression in wheat including signal transduction pathways and their components using wheat and or *A. brasilense* mutants.

**Acknowledgements.** Authors wish to thank Dr. Michael Rothballer from AMP Research Unit Microbe-Plant Interactions for providing the bacterial strains. Financial support by the University of Isfahan is gratefully acknowledged.

## REFERENCES

- [1] Agarwal, P. K., Agarwal, P., Reddy, M., Sopory, S. K. (2006): Role of DREB transcription factors in abiotic and biotic stress tolerance in plants. – *Plant cell reports* 25: 1263-1274.
- [2] Agarwal, P., Agarwal, P. K., Joshi, A. J., Sopory, S. K., Reddy, M. K. (2010): Overexpression of *PgDREB2A* transcription factor enhances abiotic stress tolerance and activates downstream stress-responsive genes. – *Molecular biology reports* 37: 11-25.
- [3] Akbari, G. A., Arab, S. M., Alikhani, H., Allakdadi, I., Arzanesh, M. (2007): Isolation and selection of indigenous *Azospirillum* spp. and the IAA of superior strains effects on wheat roots. – *World Journal of Agricultural Sciences* 3: 523-529.
- [4] Amooaghaie, R., Mostajeran, A., Emtiazi, G. (2004): The effect of compatible and incompatible *Azospirillum brasilense* strains on proton efflux of intact wheat roots. – *Plant and Soil* 243: 155-160.
- [5] Baker, S. S., Wilhelm, K. S., Thomashow, M. F. (1994): The 5'-region of *Arabidopsis thaliana* *cor15a* has cis-acting elements that confer cold-, drought-and ABA-regulated gene expression. – *Plant molecular biology* 24: 701-713.
- [6] Baldani, V. L., Baldani, J. I., Döbereiner, J. (1983): Effects of *Azospirillum* inoculation on root infection and nitrogen incorporation in wheat. – *Canadian Journal of Microbiology* 29: 924-929.
- [7] Baldani, V., Alvarez, M. D. B., Baldani, J., Döbereiner, J. (1986): Establishment of inoculated *Azospirillum* spp. in the rhizosphere and in roots of field grown wheat and sorghum. – *Plant and Soil* 90: 35-46.

- [8] Baldani, V., Baldani, J., Döbereiner, J. (1987): Inoculation of field-grown wheat (*Triticum aestivum*) with *Azospirillum* spp. in Brazil. – *Biology and fertility of soils* 4: 37-40.
- [9] Barnawal, D., Bharti, N., Pandey, S. S., Pandey, A., Chanotiya, C. S., Kalra, A. (2017): Plant growth-promoting rhizobacteria enhance wheat salt and drought stress tolerance by altering endogenous phytohormone levels and *TaCTR1/TaDREB2* expression. – *Physiologia plantarum* 161: 502-514.
- [10] Bashan, Y., Holguin, G. (1997): *Azospirillum*–plant relationships: environmental and physiological advances (1990–1996). – *Canadian Journal of Microbiology* 43: 103-121.
- [11] Beaudoin-Egan, L. D., Thorpe, T. A. (1985): Tyrosine and phenylalanine ammonialyase activities during shoot inhibition in tobacco callus cultures. – *Plant Physiology* 78: 438-441.
- [12] Bermner, E., Jenze, H., Gilbertson, C. (1995): Evidence against associative N<sub>2</sub>- fixation as a significant source of N in long-term Wheat plots. – *Plant and Soil* 175: 13-19.
- [13] Bradford, M. M. (1976): A rapid and sensitive method for the quantitation of microgram quantities of protein utilizing the principle of protein dye binding. – *Analytical Biochemistry* 72: 248-253.
- [14] Bray, E. (2000): Response to abiotic stress. – In: Bailey-Serres, J., Weretilnyk, E. (eds.) *Biochemistry and molecular biology of plants*. USA: Rockville: American Society of Plant Physiologists.
- [15] Butt, H. J., Graf, K., Kappl, M. (2006): *Physics and chemistry of interfaces*. – John Wiley & Sons.
- [16] Casimiro, I., Beeckman, T., Graham, N., Bhalerao, R., Zhang, H., Casero, P., Sandberg, G., Bennett, M. J. (2003): Dissecting *Arabidopsis* lateral root development. – *Trends in plant science* 8: 165-171.
- [17] Cohen, A. C., Bottini, R., Piccoli, P. N. (2008): *Azospirillum brasilense* Sp245 produces ABA in chemically-defined culture medium and increases ABA content in *Arabidopsis* plants. – *Plant Growth Regulation* 54: 97-103.
- [18] Correa-Aragunde, N., Graziano, M., Lamattina, L. (2004): Nitric oxide plays a central role in determining lateral root development in tomato. – *Planta* 218: 900-905.
- [19] Díaz-Zorita, M., Fernández-Canigia, M. V. (2009): Field performance of a liquid formulation of *Azospirillum brasilense* on dryland wheat productivity. – *European journal of soil biology* 45: 3-11.
- [20] Dobbelaere, S., Vanderleyden, J., Okon, Y. (2003): Plant growth-promoting effects of diazotrophs in the rhizosphere. – *Critical reviews in plant sciences* 22: 107-149.
- [21] Döbereiner, J., Day, J. M. (1976): Association symbiosis in tropical grasses: Characterization of microorganisms and dinitrogen fixing sites. – In: Newton, W. E., Nyman, C. J. (eds.) *Proceedings of the first International Symposium on Nitrogen Fixation*. Pullman: Washington State University Press.
- [22] Egawa, C., Kobayashi, F., Ishibashi, M., Nakamura, T., Nakamura, C., Takumi, S. (2006): Differential regulation of transcript accumulation and alternative splicing of a *DREB2* homolog under abiotic stress conditions in common wheat. – *Genes & genetic systems* 81: 77-91.
- [23] El-Khawas, H., Adachi, K. (1999): Identification and quantification of auxins in culture media of *Azospirillum* and *Klebsiella* and their effect on rice roots. – *Biology and Fertility of Soils* 28: 377-381.
- [24] El-Samad, A., Hetta, A. (2005): Effect of molybdenum on nitrogenase and nitrate reductase activities of wheat inoculated with *Azospirillum brasilense* grown under drought stress. – *General Applied Plant Physiology* 31: 43-54.
- [25] Estabrook, E. M., Sengupta-Gopalan, C. (1991): Differential expression of phenylalanine ammonia-lyase and chalcone synthase during soybean nodule development. – *The Plant Cell* 3: 299-308.



- [26] Fujita, Y., Fujita, M., Shinozaki, K., Yamaguchi-Shinozaki, K. (2011): ABA-mediated transcriptional regulation in response to osmotic stress in plants. – *Journal of plant research* 124: 509-525.
- [27] Gachomo, E. W., Kefela, T., Houngnandan, P., Baba-Moussa, L., Kotchoni, S. O. (2014): *Bradyrhizobium japonicum* IRAT FA3 increases biomass, yield and drought tolerance in plants. – *Journal of Natural Biology* 1: 12-23.
- [28] Gamas, P., De Billy, F., Truchet, G. (1998): Symbiosis-specific expression of two *Medicago truncatula* nodulin genes, *MtN1* and *MtN13*, encoding products homologous to plant defense proteins. – *Molecular Plant-Microbe Interactions* 11: 393-403.
- [29] Gupta, K., Agarwal, P. K., Reddy, M., Jha, B. (2010): SbDREB2A, an A-2 type DREB transcription factor from extreme halophyte *Salicornia brachiata* confers abiotic stress tolerance in *Escherichia coli*. – *Plant Cell Reports* 29: 1131-1137.
- [30] Hamdia, M. a. E.-S., Shaddad, M., Doaa, M. M. (2004): Mechanisms of salt tolerance and interactive effects of *Azospirillum brasilense* inoculation on maize cultivars grown under salt stress conditions. – *Plant Growth Regulation* 44: 165-174.
- [31] Hartmann, A., Zimmer, W., Okon, Y. (1994): *Physiology of Azospirillum*.
- [32] Hartmann, A., Baldani, J. I. (2006): The genus *Azospirillum*. – In: Dworkin, M., Falkow, S., Rosenberg, E., Schleifer, K. H., Stackebrandt, E. (eds.) *The prokaryotes Proteobacteria: alpha and beta subclasses* New York: Springer.
- [33] Hungria, M., Campo, R. J., Souza, E. M., Pedrosa, F. O. (2010): Inoculation with selected strains of *Azospirillum brasilense* and *A. lipoferum* improves yields of maize and wheat in Brazil. – *Plant and Soil* 331: 413-425.
- [34] Jain, D. K., Patriquin, D. G. (1984): Root Hair Deformation, Bacterial Attachment, and Plant Growth in Wheat-*Azospirillum* Associations. – *Applied and Environmental Microbiology* 48: 1208.
- [35] Jakab, G., Ton, J., Flors, V., Zimmerli, L., Métraux, J., Mauch-Mani, B. (2005): Enhancing *Arabidopsis* salt and drought stress tolerance by chemical priming for its abscisic acid responses. – *Plant physiology* 139: 267-274.
- [36] Jia, H., Zhang, S., Ruan, M., Wang, Y., Wang, C. (2012): Analysis and application of RD29 genes in abiotic stress response. – *Acta Physiologia Plantarum* 34: 1239-1250.
- [37] Kaur, L., Zhawar, V. K. (2015): Phenolic parameters under exogenous ABA, water stress, salt stress in two wheat cultivars varying in drought tolerance. – *Indian Journal of Plant Physiology* 20: 151-156.
- [38] Kramer, P. J., Boyer, J. S. (1995): *Water relations of plants and soils*. – Academic press.
- [39] Krieg, N., Dobreiner, J. (1984): Genus *Azospirillum*. – In: Nr, K., Jg, H. (eds.) *Bergey's Manual of Systematic Bacteriology*. London: Williams and Wilkins, Baltimore, MD.
- [40] Lawton, M. A., Dixon, R. A., Hahlbrock, K., Lamb, C. J. (1983): Elicitor induction of mRNA activity. Rapid effects of elicitor on phenylalanine ammonia-lyase and chalcone synthase mRNA activities in bean cells. – *European Journal of Biochemistry* 130(1): 131-139.
- [41] Liu, H.-S., Li, F. M., Xu, H. (2004): Deficiency of water can enhance root respiration rate of drought-sensitive but not drought-tolerant spring wheat. – *Agricultural water management* 64: 41-48.
- [42] Liu, F., Andersen, M. N., Jacobsen, S. E., Jensen, C. R. (2005): Stomatal control and water use efficiency of soybean (*Glycine max* L. Merr.) during progressive soil drying. – *Environmental and Experimental Botany* 54: 33-40.
- [43] Livak, K. J., Schmittgen, T. D. (2001): Analysis of relative gene expression data using real-time quantitative PCR and the  $2^{-\Delta\Delta CT}$  method. – *Methods* 25: 402-408.
- [44] Maksimov, I., Veselova, S., Nuzhnaya, T., Sarvarova, E., Khairullin, R. (2015): Plant growth-promoting bacteria in regulation of plant resistance to stress factors. – *Russian Journal of Plant Physiology* 62: 715-726.
- [45] Mangmang, J. S., Deaker, R., Rogers, G. (2015): *Azospirillum brasilense* enhances recycling of fish effluent to support growth of tomato seedlings. – *Horticulturae* 1: 14-26.

- [46] Mcfarland, J. (1907): The nephelometer: an instrument for estimating the number of bacteria in suspensions used for calculating the opsonic index and for vaccines. – Journal of the American Medical Association 49: 1176-1178.
- [47] Michiels, K., Croes, C. L., Vanderleyden, J. (1991): Two different modes of attachment of *Azospirillum brasilense* Sp7 to Wheat roots. – Journal of General Microbiology 137: 2241-2246.
- [48] Murashige, T., Skoog, F. (1962): A revised medium for rapid growth and bio assays with tobacco tissue cultures. – Physiologia plantarum 15: 473-497.
- [49] Nabti, E., Sahnoune, M., Ghoul, M., Fischer, D., Hofmann, A., Rothballer, M., Schmid, M., Hartmann, A. (2010): Restoration of growth of durum wheat (*Triticum durum* var. waha) under saline conditions due to inoculation with the rhizosphere bacterium *Azospirillum brasilense* NH and extracts of the marine alga *Ulva lactuca*. – Journal of Plant Growth Regulation 29: 6-22.
- [50] Nemat Alla, M. M., Younis, M. E., El-Shihaby, O. A., El-Bastawisy, Z. M. (2002): Kinetin regulation of growth and secondary metabolism in waterlogging and salinity treated *Vigna sinensis* and *Zea mays*. – Acta Physiologia Plantarum 24: 19-27.
- [51] Okon, Y., Albrecht, S. L., Burris, R. (1977): Methods for growing *Spirillum lipoferum* and for counting it in pure culture and in association with plants. – Applied and Environmental Microbiology 33: 85-88.
- [52] Pakdaman, N., Mostajeran, A., Hojati, Z. (2014): Phosphate concentration alters the effective bacterial quorum in the symbiosis of *Medicago truncatula*-*Sinorhizobium meliloti*. – Symbiosis 62: 151-155.
- [53] Pereyra, M., Zalazar, C., Barassi, C. (2006): Root phospholipids in *Azospirillum*-inoculated wheat seedlings exposed to water stress. – Plant Physiology and Biochemistry 44: 873-879.
- [54] Perrig, D., Boiero, M. L., Masciarelli, O., Penna, C., Ruiz, O. A., Cassan, F. D., Luna, M. V. (2007): Plant-growth-promoting compounds produced by two agronomically important strains of *Azospirillum brasilense* and implications for inoculant formulation. – Applied Microbiology and Biotechnology 75: 1143-1150.
- [55] Redmann, R. (1974): Osmotic and specific ion effects on the germination of alfalfa. – Canadian Journal of Botany 52: 803-808.
- [56] Rock, C. D. (2000): Pathways to abscisic acid-regulated gene expression. – New Phytologist 148: 357-396.
- [57] Sakuma, Y., Liu, Q., Dubouzet, J. G., Abe, H., Shinozaki, K., Yamaguchi-Shinozaki, K. (2002): DNA-binding specificity of the ERF/AP2 domain of *Arabidopsis* DREBs, transcription factors involved in dehydration-and cold-inducible gene expression. – Biochemical and biophysical research communications 290: 998-1009.
- [58] Santos, R., Hérouart, D., Sigaud, S., Touati, D., Puppo, A. (2001): Oxidative burst in alfalfa-*Sinorhizobium meliloti* symbiotic interaction. – Molecular plant-microbe interactions 14: 86-89.
- [59] Saubidet, M. I., Fatta, N., Barneix, A. J. (2002): The effect of inoculation with *Azospirillum brasilense* on growth and nitrogen utilization by wheat plants. – Plant and soil 245: 215-222.
- [60] Serraj, R., Sinclair, T. (2002): Osmolyte accumulation: can it really help increase crop yield under drought conditions? – Plant, cell and environment 25: 333-341.
- [61] Shen, Y. G., Zhang, W. K., He, S. J., Zhang, J. S., Liu, Q., Chen, S. Y. (2003): An EREBP/AP2-type protein in *Triticum aestivum* was a DRE binding transcription factor induced by cold, dehydration and ABA stress. – Theoretical and Applied Genetics 106: 923-930.
- [62] Shiqing, G., Huijun, X., Xianguo, C., Ming, C., Zhaoshi, X., Liancheng, L., Xingguo, Y., Lipu, D., Xiaoyan, H., Youzhi, M. (2005): Improvement of wheat drought and salt tolerance by expression of a stress-inducible transcription factor GmDREB of soybean (*Glycine max*). – Chinese Science Bulletin 50: 2714-2723.

- [63] Spaepen, S., Vanderleyden, J., Okon, Y. (2009): Plant growth-promoting actions of rhizobacteria. – *Advances in botanical research* 51: 283-320.
- [64] Sriskandarajah, S., Kennedy, I. R., Yu, D., Tchan, Y. T. (1993): Effects of plant growth regulators on acetylene-reducing associations between *Azospirillum brasilense* and wheat. – *Plant and soil* 153: 165-178.
- [65] Stein, N., Feuillet, C., Wicker, T., Schlagenhauf, E., Keller, B. (2000): Subgenome chromosome walking in wheat: a 450-kb physical contig in *Triticum monococcum* L. spans the Lr10 resistance locus in hexaploid wheat (*Triticum aestivum* L.). – *Proceedings of the National Academy of Sciences* 97: 13436-13441.
- [66] Stets, M., Alqueres, S., Souza, E., Schmid, M., Hartmann, A., Cruz, L. (2015): Quantification of *Azospirillum brasilense* FP2 Bacteria in Wheat Roots by Strain-Specific Quantitative PCR. – *Applied and environmental microbiology* 81: 6700-6709.
- [67] Sumner, M. (1990): Crop responses to *Azospirillum* inoculation. – *Advances in Soil Science* 12: 53-123.
- [68] Sweet, H. C., Bolton, W. E. (1979): The surface decontamination of seeds to produce axenic seedlings. – *American Journal of Botany* 66: 692-698.
- [69] Taylor, I. B., Burbidge, A., Thompson, A. J. (2000): Control of abscisic acid synthesis. – *Journal of Experimental Botany* 51: 1563-1574.
- [70] Tekle, A., Alemu, M. (2016): Drought Tolerance Mechanisms in Field Crops. – *World Journal of Biology and Medical Sciences* 3: 19-39.
- [71] Thomson, W. (1872): On the equilibrium of vapour at a curved surface of liquid. – *Proceedings of the Royal Society of Edinburgh* 7: 63-68.
- [72] Vasse, J., De Billy, F., Truchet, G. (1993): Abortion of infection during the *Rhizobium meliloti*—alfalfa symbiotic interaction is accompanied by a hypersensitive reaction. – *The Plant Journal* 4: 555-566.
- [73] Veresoglou, S. D., Menexes, G. (2010): Impact of inoculation with *Azospirillum spp.* on growth properties and seed yield of wheat: a meta-analysis of studies in the ISI Web of Science from 1981 to 2008. – *Plant and Soil* 337: 469-480.
- [74] Yamaguchi-Shinozaki, K., Shinozaki, K. (2005): Improving drought and cold-stress tolerance in transgenic rice. *Rice is life: scientific perspectives for the 21st century.* – *Proceedings of the World Rice Research Conference held in Tsukuba, Japan, 4-7 November 2004, 2005.* International Rice Research Institute (IRRI), 94-97.
- [75] Yang, J., Kloepper, J. W., Ryu, C.-M. (2009): Rhizosphere bacteria help plants tolerate abiotic stress. – *Trends in plant science* 14: 1-4.

## APPENDIX

**Appendix 1.** The averages of ranking numbers in the associated pairs between wheat (*Triticum aestivum* L.) cultivars (Omid, Roshan, Sardari, Shahpasand, Shoaleh and Tabasi) and two strains of *A. brasilense* (Sp7, Sp245) at optimum water content (80% FC) and drought condition (40% FC) based on growth and biochemical parameters. At 80% FC, the minimum and maximum numbers show the most compatible (homologous) and most non-compatible (heterologous) association pairs, respectively; At 40% FC, the minimum and maximum numbers show the most tolerant and most sensitive association pairs to drought, respectively; At 40% FC, Tabasi cultivar is not included due to lack of its successful association with both bacterial strains under optimum water condition (NI: non-inoculated)

Pairs		Ranking number	
Wheat cultivars	<i>A. brasilense</i> strains	80% FC	40% FC
Omid	Non-inoculated	8.2	8.5
	Sp7	6.3	7.2
	Sp245	4.5	5
Sardari	Non-inoculated	12.2	10
	Sp7	9	7.3
	Sp245	7.7	5.5
Roshan	Non-inoculated	6.5	4
	Sp7	3.3	2.7
	Sp245	<b>3 (Minimum)</b>	<b>1 (Minimum)</b>
Shoaleh	Non-inoculated	13.8	15.0
	Sp7	10.3	14
	Sp245	7.5	11.2
Tabasi	Non-inoculated	12.5	-
	Sp7	10.5	-
	Sp245	10.5	-
Shahpasand	Non-inoculated	12.2	10.8
	Sp7	<b>11 (Maximum)</b>	<b>16 (Maximum)</b>
	Sp245	8	11.7