

Effect of Hydatid Cyst Fluid Antigens on Induction of Apoptosis on Breast Cancer Cells

Abstract

Background: Hydatid cyst is a zoonotic and parasitic disease with worldwide distribution. Anticancer effects of hydatid cyst have been shown in cell culture experiments and animal model investigations. The mechanism of anti-cancer effects of hydatid cyst fluid has not been clearly elucidated, and the induction of apoptosis may have a role in this regard. Hence, in this study, the effect of hydatid cyst fluid (HCF) on the induction of apoptosis on mouse breast cancer (4T1) cell line in cell culture medium has been investigated. **Materials and Methods:** *Echinococcus granulosus* HCF antigens including Antigen B (AgB), glycolipid, glycoprotein, and 78 KDa fractions were prepared. Breast cancer cell line (4T1) was cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum and appropriate antibiotics. Apoptosis was determined by flow cytometry using the annexin V-fluorescein isothiocyanate apoptosis kit. **Results:** The 78 KDa and glycoprotein fractions induced more than 40% apoptosis. HCF and glycolipid antigens induced 39% and 34% apoptosis, respectively. However, less apoptosis observed after treatment with AgB fraction. **Conclusion:** Hydatid cyst antigens especially the 78 KDa and glycoprotein fractions induced apoptosis on mouse breast cancer cells.

Keywords: 4T1 cells, apoptosis, cancer, hydatid cyst

Introduction

Hydatid cyst is the larval stage of the tapeworm *Echinococcus granulosus* (*E. granulosus*) which is in human and livestock viscera. The cyst is outwardly covered with a carbohydrate-rich material termed laminated layer. The hydatid cyst fluid (HCF) of *E. granulosus* is a complex of biological mixture of glycoprotein and glycolipid, carbohydrates, cyclophilin, and ferritin.^[1]

Apoptosis is one of the mechanisms that provides a suitable situation for survival of the cyst by inducing apoptosis in the host immune cells. Several factors play a role in apoptosis, but the caspase enzymes have the prominent role.^[2,3] In another investigation, cytotoxic features of HCF have been shown.^[4,5] On the other hand, in many studies, the ability of HCF to suppress cancer growth has been well documented both in cell culture experiments^[6,7] and in experimental animal model investigations.^[8,9]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Breast cancer is the second leading cause of cancer death among women and is the most common malignancy in women accounting for 1.2 million new cases annually.^[10] First-line treatment options are surgery, radiation, hormonal therapy, and chemotherapy. Moreover, nonselective cytotoxicity of chemotherapeutic agents causes adverse effects. Furthermore, drug resistance makes a weak prognosis in the treatment process.^[11,12] Therefore, searching for effective anti-cancer agents with higher efficiency and fewer side effects is important. Hence, in this study, apoptotic and necrotic features of HCF antigens on mouse breast cancer (4T1) cells has been investigated.

Materials and Methods

HCF was collected from infected sheep lungs and livers at Fasaran slaughterhouse in Isfahan, Iran. From sheep hydatid cyst, the hydatid fluid was aspirated using a syringe and needle. After that with the observation of the protoscolices, the

How to cite this article: Daneshpour S, Kefayat AH, Mofid MR, Rostami Rad S, Yousofi Darani H. Effect of hydatid cyst fluid antigens on induction of apoptosis on breast cancer cells. *Adv Biomed Res* 2019;8:27.

Received: October, 2018. **Accepted:** February, 2019.

Shima Daneshpour,
Amir Hossein
Kefayat¹,
Mohammad Reza
Mofid²,
Shahla Rostami
Rad³,
Hossein Yousofi
Darani

From the Department of Parasitology and Mycology, Medical School, Isfahan University of Medical Sciences, ¹Cancer Prevention Research Centre, Isfahan University of Medical Sciences, ²Department of Biochemistry, Bioinformatics Research Center, School of Pharmacy and Pharmaceutical Sciences, Isfahan University of Medical Sciences, Isfahan, ³Department of Medicine, Najafabad Branch, Islamic Azad University, Najafabad, Iran

Address for correspondence:
Prof. Hossein Yousofi Darani,
Department of Medical
Parasitology and Mycology,
Isfahan University of Medical
Sciences, Isfahan, Iran.
E-mail: yousofidarani@gmail.com

Access this article online

Website: www.advbiores.net

DOI: 10.4103/abr.abr_220_18

Quick Response Code:



aspirated cyst fluids were centrifuged at $2000 \times g$, for 2 min and the supernatant was air dried and stored at -20°C as crud hydatid fluid antigen.

A sample of CHF with 2 mg/ml concentration was purified by gel filtration chromatography in a column on Sephadex G 200 (1.5 cm diameter and 60 cm in length) in phosphate buffered saline (PBS) (pH 7.2), at a flow rate of 20 ml/hour. The elutes were collected and pooled together. The protein concentration determined by bicinchoninic acid protein kit.^[13] The purified fractions were then analyzed in sodium dodecyl sulfate-polyacrylamide gel electrophoresis (SDS-PAGE), (12.5%) according to Laemmli method^[14] with some modification. The gel was then stained with silver stain. Molecular weights were determined by standard protein markers (PMW-M, Bangalore Genei).

Preparation of antigen B of hydatid cyst fluid

The antigen B (AgB) of HCF fraction was prepared according to the previously described method.^[15] Briefly, 50 mL tubes containing HCF were centrifuged at 1500 g for 15 min. The supernatant was then dialyzed against 0.005 M acetate buffer (pH = 5) overnight at 4°C and centrifuged at 5000 g for 30 min. Then, the precipitate was dissolved in 10 ml of 2M phosphate buffer (pH = 8). The preparation was saturated with 5 mL ammonium sulfate 40% and mixed for 1 h and centrifuged at 5000 g for 15 min. The supernatant was incubated in water bath for 15 min and centrifuged at 5000 for 30 min. Finally, the supernatant containing the Ag B was collected and stored at -70°C until use.

Preparation of glycoprotein and glycolipid antigens of hydatid cyst fluid

The glycoprotein and glycolipid of HCF were prepared by chloroform-methanol extraction. Distilled water, methanol, and chloroform were added to the crude HCF with 3:4:1:1 ratio, respectively and centrifuged for about 5 min at 5000 g. The pellet was then resuspended in 4 ml methanol and centrifuged at 5000 g for 20 min. After discarding the supernatant and resuspension of

the pellet, the lower section of the tube was obtained as the glycoprotein fraction and the supernatant as the glycolipid fraction.

Cell culture

Breast cancer cell line (4T1) was obtained from the National Cell Bank of Iran (Pasteur Institute, Tehran, Iran). Cells were cultured in RPMI-1640 medium supplemented with 10% fetal bovine serum, 100 ng/mL of streptomycin and 100 units/mL of penicillin and kept at 37°C in a 5% CO_2 incubator.

Apoptosis assay

By using the annexin V-fluorescein isothiocyanate (V-FITC) apoptosis kit (eBioscience, USA) apoptotic rate was determined by flow cytometry.^[16] The rate of apoptotic cells was estimated by green fluorescence emitted by annexin V-FITC bound to phosphatidylserine exposed to the outer leaflet of the apoptotic cell membrane. After extensive washing in PBS, 1×10^6 cells were rinsed with HEPES buffer, re-suspended in the same buffer, and incubated at room temperature for 5–15 min in the dark and then annexin V-FITC (5 μl) was added. To detect necrotic cells, PI was then added to the cell suspension before the analyzing by a FACS equipment (BD Biosciences, San Jose, CA, USA).

Statistical analysis

Data were expressed as mean \pm standard deviation from three independent experiments. Statistical analysis was performed using ANOVA and Student's t-tests, and the differences were considered statistically significant at $P < 0.05$.

Results

Results of gel filtration chromatography

Results of gel filtration chromatography of HCF showed two peaks [Figure 1]. In the first one (peak 1), the protein concentration was 600 $\mu\text{g}/\text{ml}$, but in the second one (peak 2) the main constituent was urea. SDS-PAGE analysis of the proteins in peak 1 showed that a band with a molecular weight of about 78 KDa [Figure 2].




Figure 1: Gel filtration chromatography of hydatid cyst fluid

Effect of different antigens of hydatid cyst fluid on induction of apoptosis on breast cancer cells

The effects of different antigens of hydatid cyst on the induction of apoptosis and necrosis were estimated following 48 h after the treatment of breast cancer cells with hydatid cyst antigens. The 78 KDa fraction of HCF induced 46.5% apoptosis. Glycoprotein fraction and crude HCF induced 41% and 39% apoptosis, respectively. Glycolipid fraction induced 34% apoptosis and 20% apoptosis observed following treatment with AgB. In the control group (nontreated cells), 17% apoptosis was also observed. However, no obvious amount of necrosis was seen following treatment of breast cancer cells with hydatid cyst antigens. The results of apoptosis and necrosis induction by different hydatid cyst antigens have been summarized in Figures 3 and 4.

Discussion

Results of this work indicated that HCF crude antigen; the 78KD, and glycoprotein fractions induced apoptosis on breast cancer cells. *In vitro* and *in vivo* studies demonstrate that some parasite has anticancer effects. In this context, it has been shown that hydatid cyst protoscolices induce cell death in WEHI 164 fibrosarcoma cells.^[6] Also, the inhibitory effect of different hydatid cyst molecules on Hela and Vero cell lines has been shown.^[7] In another studies, HCF reduces the growth of melanoma tumor in C57/black mice.^[8] Furthermore, antitumor activity of human HCF in a murine model of colon cancer has been shown.^[9] Karadayi *et al.* show that hydatid disease has protective effects against lung cancer.^[17]

How hydatid cyst-induced cell death and associated signal pathway of apoptosis are not clear. It was previously shown that hydatid fluid toxins have fatal effects on mouse peritoneal macrophages. Thus, apoptosis pathways

may be activated after exposure to hydatid fluid by secreted toxins of the protoscolices. In an investigation, it has been shown that hydatid cyst induced apoptosis through increased Bcl-2 family levels.^[18] Cytotoxic features of *E. granulosus* has previously been shown. In an investigation, it has been shown that in lymphocytes that were treated by fertile hydatid fluid apoptosis rate was significantly higher in comparison to nontreated cells. In the treated lymphocytes, expression of the Bax gene as a pro-apoptotic molecule was increased and the expression level of Bcl-2 mRNA as an anti-apoptotic molecule was decreased. The activity of caspase-3 was also significantly higher in the hydatid cyst-treated group.^[19] All these support this concept that antigens of protoscolices of *E. granulosus* HCF induce apoptosis in the lymphocytes of hosts. However, in a research work done by Janssen *et al.*, it was demonstrated that some toxic substances of the protoscolices, may reduce the viability of the macrophages *in vitro*.^[20] Nono *et al.* demonstrate that excretory/secretory secretions of *Echinococcus multilocularis* (*E. multilocularis*) cyst induce apoptosis in dendritic cells *in vitro*.^[21] In another work, Emp53 was identified from *E. multilocularis* and was shown that this molecule has apoptosis and the oxidative stress effects.^[22] In conclusion, our results confirmed that some antigens of hydatid cyst fluid induce apoptosis in 4T1 cells. Hence, HCF may be a curative material to induce apoptosis in cancer cells. However, the molecular mechanisms associated with the anti-apoptosis of hydatid antigens have not been clearly elucidated. More researches are recommended to understand molecular mechanisms of apoptosis induced by the parasite.

Financial support and sponsorship

This work was supported by grant from Isfahan University of Medical Sciences, Isfahan, Iran.

Conflicts of interest

There are no conflicts of interest.




Figure 2: Sodium dodecyl sulfate-polyacrylamide gel electrophoresis of protein in Peak 1 of hydatid cyst fluid




Figure 3: Effects of different antigens of hydatid cyst the 78 KD fraction (a), glycoprotein (b), crud hydatid cyst (c), glycolipid (d), antigen B (e), and control (f) on apoptosis and necrosis of 4T1 cells. The values of necrosis and apoptosis were shown in percentage (%)




Figure 4: Flow cytometry analysis of effects of different antigens of hydatid cyst the 78 KD fraction (a), glycoprotein (b), crud hydatid cyst (c), glycolipid (d), antigen B (e), and control (f) on apoptosis and necrosis of 4T1 cells

References

1. Aziz A, Zhang W, Li J, Loukas A, McManus DP, Mulvenna J, *et al.* Proteomic characterisation of *Echinococcus granulosus* hydatid cyst fluid from sheep, cattle and humans. *J Proteomics* 2011;74:1560-72.
2. Cabrera G, Cabrejos ME, Morassutti AL, Cabezón C, Orellana J, Hellman U, *et al.* DNA damage, RAD9 and fertility/infertility of *Echinococcus granulosus* hydatid cysts. *J Cell Physiol* 2008;216:498-506.
3. Paredes R, Jiménez V, Cabrera G, Iragüen D, Galanti N. Apoptosis as a possible mechanism of infertility in *Echinococcus granulosus* hydatid cysts. *J Cell Biochem* 2007;100:1200-9.
4. MacIntyre AR, Dixon JB, Green JR. Growth kinetics of leukocyte cell lines cultured with hydatid fluid of *Echinococcus granulosus* equinus. *Parasite Immunol* 2000;22:651-7.
5. Macintyre AR, Dixon JB, Green JR. Mitosis and differentiation in T-cells under cytotoxic action of *Echinococcus granulosus* hydatid fluid. *Vet Parasitol* 2001;96:277-89.
6. Yousofi Darani H, Soozangar N, Khorami S, Taji F, Yousofi M, Shirzad H, *et al.* Hydatid cyst protoscolices induce cell death in WEHI-164 fibrosarcoma cells and inhibit the proliferation of baby hamster kidney fibroblasts *in vitro*. *J Parasitol Res* 2012;2012:304183.
7. Aref N, Shirzad H, Yousefi M, Darani H. Effect of different hydatid cyst molecules on hela and vero cell lines growth *in vitro*. *J Immunodeficient Disor* 2012;2:3.
8. Chookami MB, Sharafi SM, Sefiddashti RR, Jafari R, Bahadoran M, Pestechian N, *et al.* Effect of two hydatid cyst antigens on the growth of melanoma cancer in C57/black mice. *J Parasit Dis* 2016;40:1170-3.
9. Berriel E, Russo S, Monin L, Festari MF, Berois N, Fernández G, *et al.* Antitumor activity of human hydatid cyst fluid in a murine model of colon cancer. *ScientificWorldJournal* 2013;2013:230176.
10. Parkin DM, Bray F, Ferlay J, Pisani P. Global cancer statistics, 2002. *CA Cancer J Clin* 2005;55:74-108.
11. Sharifi S, Barar J, Hejazi MS, Samadi N. Roles of the Bcl-2/Bax ratio, caspase-8 and 9 in resistance of breast cancer cells to paclitaxel. *Asian Pac J Cancer Prev* 2014;15:8617-22.
12. Ghanbari P, Mohseni M, Tabasinezhad M, Yousefi B, Saei AA, Sharifi S, *et al.* Inhibition of survivin restores the sensitivity of breast cancer cells to docetaxel and vinblastine. *Appl Biochem Biotechnol* 2014;174:667-81.
13. He F. BCA (Bicinchoninic Acid) Protein Assay. *Bio Protocol Biol* 2011;101:e44.
14. Laemmli UK. Cleavage of structural proteins during the assembly of the head of bacteriophage T4. *Nature* 1970;227:680-5.
15. Shirazi S, Madani R, Hoghooghi Rad N, Bahadori R. Isolation and purification of *Echinococcus granulosus* antigen B from hydatid cyst fluid using three different methods. *Arch Razi Instit* 2016;71:103-8.
16. Rieger AM, Nelson KL, Konowalchuk JD, Barreda DR. Modified annexin V/propidium iodide apoptosis assay for accurate assessment of cell death. *J Vis Exp* 2011. pii: 2597.
17. Karadayi S, Arslan S, Sumer Z, Turan M, Sumer H, Karadayi K, *et al.* Does hydatid disease have protective effects against lung cancer? *Mol Biol Rep* 2013;40:4701-4.
18. Janssen D, De Rycke PH, Osuna A. Dose-dependent effects of hydatid fluid toxins from *Echinococcus granulosus* on mouse peritoneal macrophages. *Folia Parasitol (Praha)* 1993;40:109-13.
19. Mokhtari Amirajdi M, Sankian M, Eftekharzadeh Mashhadi I,

- Varasteh A, Vahedi F, Sadrizadeh A, *et al.* Apoptosis of human lymphocytes after exposure to hydatid fluid. *Iran J Parasitol* 2011;6:9-16.
20. Janssen D, Osuna A, Lazuen J, de Rycke PH. Comparative cytotoxicity of secondary hydatid cysts, protoscoleces, and *in vitro* developed microcysts of *Echinococcus granulosus*. *J Helminthol* 1992;66:124-31.
21. Nono JK, Pletinckx K, Lutz MB, Brehm K. Excretory/secretory-products of *Echinococcus multilocularis* larvae induce apoptosis and tolerogenic properties in dendritic cells *in vitro*. *PLoS Negl Trop Dis* 2012;6:e1516.
22. Cheng Z, Zhu S, Wang L, Liu F, Tian H, Pengsakul T, *et al.* Identification and characterisation of emp53, the homologue of human tumor suppressor p53, from *Echinococcus multilocularis*: Its role in apoptosis and the oxidative stress response. *Int J Parasitol* 2015;45:517-26.

To,
The Editor

Covering Letter

Submission of Manuscript for publication

Dear Sir,

We intend to publish an article entitled

in your journal.

On behalf of all the contributors I will act and guarantor and will correspond with the journal from this point onward.

Prior presentation of the data reported in this manuscript:

Organisation

Place

Date

We have done sufficient work in the field to justify authorship for this manuscript.

We hereby transfer, assign, or otherwise convey all copyright ownership, including any and all rights incidental thereto, exclusively to the journal, in the event that such work is published by the journal.

Thank you,

Yours' sincerely,

Name of corresponding contributor

Signature

Title of the manuscript:

Title Page

Type of manuscript:

Running title:

Contributors:

	First name	Middle name initial	Last name	Highest academic degree	Names of departments and institutions (including city and state)	Email addresses
1						
2						
3						
4						
5						
6						

Corresponding Author:

Name:

Address:

Phone numbers:

Facsimile numbers:

E-mail address:

Total number of pages:

Total number of tables:

Total number of figures:

Total number of supplementary files:

Word counts: For abstract:

For the text:

Acknowledgement:

Conflict of interest:

Financial Support:

Contribution details (to be ticked marked as applicable):

Contributors' form

	Contributor 1	Contributor 2	Contributor 3	Contributor 4	Contributor 5	Contributor 6
Concepts						
Design						
Definition of intellectual content						
Literature search						
Clinical studies						
Experimental studies						
Data acquisition						
Data analysis						
Statistical analysis						
Manuscript preparation						
Manuscript editing						
Manuscript review						
Guarantor						