

Essential hypertension in children, a growing worldwide problem

Mohammadreza Sabri*¹, Alaleh Gheissari*², Marjan Mansourian³, Noushin Mohammadifard⁴, Nizal Sarrafzadegan⁵

¹Pediatric Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran,

²Department of Pediatrics, School of Medicine, Isfahan University of Medical Sciences, Isfahan, Iran, ³Department of Epidemiology and

Biostatistics, School of Health, Isfahan University of Medical Sciences, Isfahan, Iran, ⁴Hypertension Research Center, Cardiovascular

Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran, ⁵Isfahan Cardiovascular Research Center, Cardiovascular Research Institute, Isfahan University of Medical Sciences, Isfahan, Iran

*They are equally contributed

Hypertension is one of the most common diseases worldwide. For many decades, it was considered as a problem related to adult population; however, its incidence in children has also been increased in recent years. Although secondary causes of hypertension are more common in children, few studies have been published focusing on the growing epidemic rate of essential hypertension in children and adolescents. Considering the importance of essential hypertension and its cardiovascular consequences, we review briefly its epidemiology and risk factors in children.

Key words: Adolescent, cardiovascular disease, child, essential hypertension, obesity

How to cite this article: Sabri M, Gheissari A, Mansourian M, Mohammadifard N, Sarrafzadegan N. Essential hypertension in children, a growing worldwide problem. *J Res Med Sci* 2019;24:109.

INTRODUCTION

Hypertension has been known not only as a common disease but also as one of the most prevalent human diseases, leading to high morbidity and mortality.^[1] About 10.4 million deaths were attributed to high systolic blood pressure as the leading risk factor of cardiovascular disease.^[2] In the past, hypertension and its harmful consequences were attributed to adult patients, and it was unusual in childhood and secondary to some disorders of the renal, endocrine, and cardiac.^[3]

One of the first reports on the prevalence of pediatric hypertension was published in 1963.^[4] Since then, many papers have been published regarding the prevalence and importance of hypertension in pediatric population.

While primary hypertension is more common in the adult population, secondary causes of

hypertension are mostly found in children. Renal and renovascular diseases have their top position among identifiable causes of hypertension in children.^[5] Glomerulonephritis, reflux nephropathy, renal artery stenosis, and coarctation of the aorta followed by endocrine disorders are main causes of secondary hypertension in the pediatric population.^[3,6,7] Thus, in this paper, we aim to review the epidemiology of primary hypertension and its determinants including gender, obesity, lifestyles, and genetic factors in children and adolescents.

EPIDEMIOLOGY

In recent decades, an increase in primary hypertension in adolescents has been reported.^[7-9] The prevalence of essential hypertension in adolescents is different among diverse ethnicities from as low as 0.3% up to approximately 21%.^[8-13]

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: reprints@medknow.com

Access this article online

Quick Response Code:



Website:

www.jmsjournal.net

DOI:

10.4103/jrms.JRMS_641_19

Address for correspondence: Prof. Alaleh Gheissari, Department of Pediatrics, School of Medicine, Isfahan University of Medical Sciences, Hezar Jerib St, Isfahan, Iran. E-mail: alaleh.gheissari@gmail.com

Received: 26-08-2019 **Revised:** 30-08-2019 **Accepted:** 03-09-2019

Various factors such as obesity, diet, gender, and academic stress have been claimed as responsible causes in increasing the prevalence of primary hypertension in this age group.^[11,13-15]

Gender

Some studies stated that essential hypertension is more common in boys than in girls. Sundar *et al.* reported a high rate of essential hypertension in boys, with a male/female ratio approximately equals to 3/1.^[19] In another survey, Mohan *et al.* showed that hypertension was more common in boys not only in urban but also in rural area of Ludhiana, India.^[16] A study by Buch *et al.* demonstrated slight increase in the prevalence of hypertension in boys aged older than 13 years compared to girls.^[17] However, the gender effect has not been supported by all similar studies.^[18]

Obesity and body mass index

Although the rising trend of obesity and mean body mass index (BMI) in high levels has been plateaued in high-income countries, it has enhanced rapidly in the south, east, and southeast of Asia.^[19] Growing rate of obesity changes the incidence of overweight-related problems such as hypertension in the world.^[11,12,20-22]

According to the results from the National Health and Nutrition Examination Survey between 1988 and 1994 and between 1999 and 2008, in parallel with increasing BMI, 3.4% and 4.4% rise in the prevalence of hypertension have been reported among boys and girls, respectively.^[23]

By various mechanisms including insulin resistance, activation of renin-angiotensin-aldosterone system (RAS), retention of salt, and consequent changes in vascular endothelial function, obesity induces hypertension.^[24]

A study on about 25,000 schoolchildren revealed a higher prevalence of hypertension in overweight and obese children in comparison to normal weight participants (17%–18% vs. 10%).^[25]

The rise of the population with hypertension has been shown not only in adolescents but even in children. Based on the electronic medical data from a cohort of 14,000 children and adolescents aged 3–18 years, a prevalence of 3.4% of hypertensive population was reported and the presence of obesity was accompanied by higher blood pressure.^[26]

Lifestyle factors

Dietary patterns, physical activity, and stress have main roles in the incidence and prevention of hypertension in children and adolescents.^[27] A diet with high salt intake, saturated and trans-fatty acids, and low consumption of fruits and vegetables, nuts, and olive oil was associated

with hypertension in both children and adults.^[28] Due to urbanization, children have more consumption of processed food and energy-dense diet,^[29] which cause obesity and consequently obesity-related hypertension in childhood.^[30] Sedentary lifestyle and poor sleep quality promote hypertension in children.^[22] Since the World Health Organization recommended to engage in moderate-to-vigorous physical activity for at least 1 h a day in childhood.^[31] Elevated blood pressure in schoolchildren and adolescents can be attributed to high stress, particularly academic stress, because of difficult curriculum in the school and educational competition.^[32]

Genetic factors

Although excess weight is the main factor responsible for increasing rate of hypertension among adolescent, genetic, and family history, susceptibility should not be ignored.^[33,34]

Gene polymorphism of RAS has been proposed as a risk factor for essential hypertension and its cardiovascular consequences.

Aldosterone synthase gene (CYP11B2) polymorphism evaluation in Japanese participants with essential hypertension showed a significant difference in the distribution of three genotypes (TT, TC, and CC) between the hypertensive and normotensive population.^[34]

A recent study in China demonstrated the higher risk of hypertension in children and adolescents with single-nucleotide polymorphism of ATP2B1 rs17249754.^[35]

In addition to RAS, endothelial nitric oxide gene polymorphism has been targeted as a possible genetic factor to contribute to developing essential hypertension. More frequency of one of the most applicable polymorphisms in the NOS3 gene (rs1799983 in exon 7) has been reported in Sudanese patients with essential hypertension than control group.^[36]

CONCLUSION

Concerning the growing rate of essential hypertension and its link to environmental, genetic, and anthropometric factors, assessing its prevalence and possible cardiovascular complications is necessary for every population and geographical area.

Financial support and sponsorship

Nil.

Conflicts of interest

There are no conflicts of interest.

REFERENCES

- Bromfield S, Muntner P. High blood pressure: the leading global burden of disease risk factor and the need for worldwide prevention programs. *Curr Hypertens Rep* 2013;15:134-6.
- GBD 2017 Risk Factor Collaborators. Global, regional, and national comparative risk assessment of 84 behavioural, environmental and occupational, and metabolic risks or clusters of risks for 195 countries and territories, 1990-2017: A systematic analysis for the Global Burden of Disease Study 2017. *Lancet* 2018;392:1923.
- Sharma S, Meyers KE, Vidi SR. Secondary forms of hypertension in children: Overview. In: Flynn J, Ingelfinger JR, Redwine K, editors. *Pediatric Hypertension*. Cham: Springer International Publishing; 2016. p. 1-20.
- Chiolerio A, Bovet P, Paradis G, Paccaud F. Has blood pressure increased in children in response to the obesity epidemic? *Pediatrics* 2007;119:544-53.
- Baracco RI, Kapur G, Mattoo T, Jain A, Valentini R, Ahmed M, *et al.* Prediction of primary vs secondary hypertension in children. *J Clin Hypertens (Greenwich)*. 2012;14:316-21.
- Charles L, Triscott J, Dobbs B. Secondary hypertension: discovering the underlying cause. *Am Fam Physician* 2017;96:453-46.
- Stein DR, Ferguson MA. Evaluation and treatment of hypertensive crises in children. *Integr Blood Press Control* 2016;9:49-58.
- Wühl E. Hypertension in childhood obesity. *Acta Paediatr* 2019;108:37-43.
- Sundar JS, Adaikalam JM, Parameswari S, Valarmarathi S, Kalpana S, Shantharam D, *et al.* Prevalence and determinants of hypertension among urban school children in the age group of 13- 17 years in, Chennai, Tamilnadu. *Epidemiol* 2013;3:3-5.
- Bell CS, Samuel JP, Samuels JA. Prevalence of hypertension in children. *Hypertension* 2019;73:148-52.
- Gupta R. Trends in hypertension epidemiology in India. *J Hum Hypertens* 2004;18:3-8.
- Sharma A, Grover N, Kaushik S, Bhardwaj R, Sankhyan N. Prevalence of hypertension among school children in Shimla. *Indian Pediatr* 2010;47:873-6.
- Rao G. Diagnosis, epidemiology, and management of hypertension in children. *Pediatrics* 2016;138. pii: e20153616.
- Ewald DR, Haldeman LA. Risk factors in adolescent hypertension. *Glob Pediatr Health* 2016;3:2333794X15625159.
- Raj M. Essential hypertension in adolescents and children: Recent advances in causative mechanisms. *Indian J Endocrinol Metab* 2011 Suppl 4:S367-73.
- Mohan B, Kumar N, Aslam N, Rangbulla A, Kumbkarni S, Sood NK, *et al.* Prevalence of sustained hypertension and obesity in urban and rural school going children in Ludhiana. *Indian Heart J* 2004;56:310-4.
- Buch N, Goyal JP, Kumar N, Parmar I, Shah VB, Charan J. Prevalence of hypertension in school going children of Surat city, Western India. *J Cardiovasc Dis Res* 2011;2:228-32.
- Papandreou DA, Stamou MA, Malindretos PB, Rousso IA, Mavromichalis I. Prevalence of hypertension and association of dietary mineral intake with blood pressure in healthy schoolchildren from Northern Greece aged 7-15 years. *Ann Nutr Metab* 2007;51:471-6.
- NCD Risk Factor Collaboration (NCD-RisC). Worldwide trends in body-mass index, underweight, overweight, and obesity from 1975 to 2016: a pooled analysis of 2416 population-based measurement studies in 128.9 million children, adolescents, and adults. *Lancet* 2017;390:2627-42.
- Din-Dzietham R, Liu Y, Bielo MV, Shamsa F. High blood pressure trends in children and adolescents in national surveys, 1963 to 2002. *Circulation* 2007;116:1488-96.
- Harris K, Benoit G, Dionne J, Feber J, Cloutier L, Zarnke K *et al.* Hypertension Canada's 2016 Canadian Hypertension Education Program Guidelines for Blood Pressure Measurement, Diagnosis, and Assessment of Risk of Pediatric Hypertension. *Can J Cardiol* 2016;32:589e597.
- Kelly RK, Magnussen CG, Sabin MA, Cheung M, Juonala M. Development of hypertension in overweight adolescents: a review. *Adolesc Health Med Ther* 2015;6:171-8.
- Rosner B, Cook NR, Daniels S, Falkner B. Childhood blood pressure trends and risk factors for high blood pressure: the NHANES experience 1988-2008. *Hypertension* 2013;62:247-54.
- Kotchen TA. Obesity-related hypertension: Epidemiology, pathophysiology, and clinical management. *Am J Hypertens* 2010;23:1170-8.
- Raj M, Sundaram KR, Paul M, Deepa AS, Kumar RK. Obesity in Indian children: Time trends and relationship with hypertension. *Natl Med J India* 2007;20:288-93.
- Hansen ML, Gunn PW, Kaelber DC. Underdiagnosis of hypertension in children and adolescents. *JAMA* 2007;298:874-9.
- Giontella A, Bonafini S, Tagetti A, Bresadola I, Minuz P, Gaudino R, *et al.* Relation between dietary habits, physical activity, and anthropometric and vascular parameters in children attending the primary school in the Verona south district. *Nutrients* 2019;11:14.
- Casas R, Castro-Barquero S, Estruch R, Sacanella E. Nutrition and Cardiovascular Health. *Int J Mol Sci* 2018;19:3988.
- World Health Organization. Commission on ending childhood obesity. Report of the Commission on Ending Childhood Obesity. Geneva, Switzerland: World Health Organization; 2016.
- Ambrosini GL. Childhood dietary patterns and later obesity: A review of the evidence. *Proc Nutr Soc* 2014;73:137-46.
- World Health Organization. Information Sheet: Global Recommendations on Physical Activity for Health 5-17 Years Old. Geneva, Switzerland: World Health Organization; 2015.
- Arun DJ, Kavinilavu R. A study of risk factors associated with hypertension among the school going children in Puducherry. *Int J Community Med Public Health* 2018;5:764-8.
- Solanki JD, Mehta HB, Shah CJ. Pulse wave analyzed cardiovascular parameters in young first degree relatives of hypertensives. *J Res Med Sci* 2018;23:72.
- Ji X, Qi H, Li DB, Liu RK, Zheng Y, Chen HL, *et al.* Associations between human aldosterone synthase CYP11B2 (-344T/C) gene polymorphism and antihypertensive response to valsartan in Chinese patients with essential hypertension. *Int J Clin Exp Med* 2015;8:1173-7.
- Xi B, Shen Y, Zhao X, Chandak GR, Cheng H, Hou D, *et al.* Association of common variants in/near six genes (ATP2B1, CSK, MTHFR, CYP17A1, STK39 and FGF5) with blood pressure/hypertension risk in Chinese children. *J Hum Hypertens* 2014;28:32-6.
- Gamil S, Erdmann J, Abdalrahman IB, Mohamed AO. Association of NOS3 gene polymorphisms with essential hypertension in Sudanese patients: a case control study. *BMC Med Gen* 2017;18:128.

© 2019. This work is published under

<https://creativecommons.org/licenses/by-nc-sa/4.0/>(the “License”).

Notwithstanding the ProQuest Terms and Conditions, you may use this content
in accordance with the terms of the License.